



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

June 23, 2010

Ms. Laura Phelps
Regulatory Affairs, Herbicides & PGRs
Bayer CropScience
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

Sencor 4 Flowable Herbic

RE: ADMINISTRATIVE LTR DATED MAY 20, 2010; ~~RADIUS HERBICIDE~~, EPA REG.
NO. 264-735; AMENDED LABEL PER THE PESTICIDE CONTAINER AND
CONTAINMENT FINAL RULE, AUGUST 16, 2006

Your amended language request per PR Notice 2007-4: Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment." is approved with the following changes in the "STORAGE AND DISPOSAL" box;

1. Under "Pesticide Disposal:" change the current wording to "Wastes resulting from the use of this product may be disposed of on site in accordance with the uses permitted on this label, or at an approved waste disposal facility."
2. Under "Container Disposal:", everywhere it appears, change "Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.", to "Once container is rinsed, offer for recycling if available or, if allowed by State or local authorities, puncture and dispose of in a sanitary landfill."


Only the amended language in the "STORAGE AND DISPOSAL" box and the paragraph labeled "Refillable Containers" directly under the box was reviewed and approved with the above comments. Therefore, only the first three pages of the label is enclosed, and Approved with Comments. These pages replace all past versions of the same content, and constitute part of the approved label. Pages 4 to the end of the label remain as approved in the last stamped approved label for this product, plus any valid notifications.

Product released for shipment after August 16, 2010 (plus any extension published as a final rule in the Federal Register), or 18 months from the date of this letter, whichever is shorter, must

incorporate the approved language with the changes above on the pesticide label (EPA Reg. No. 264-852).

Please contact Phil Errico at 703-305-6663/ errico.philip@epa.gov for any assistance in this matter.

Regards,



James A. Tompkins PM 25
Herbicide Branch/Registration Division 7505P

Sencor[®] 4 Flowable Herbicide

For control of certain grasses and broadleaf weeds on alfalfa, asparagus, barley (spring and winter), carrots, field corn, garbanzo beans (chick peas), lentils, peas, potatoes, sainfoin, soybeans, sugarcane, tomatoes, winter wheat.

| | |
|---|------|
| ACTIVE INGREDIENT: | |
| Metribuzin: 4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one..... | 41% |
| INERT INGREDIENTS: | |
| Contains 4 lb of 4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one per gallon. | 100% |

EPA Reg. No. 264-735 EPA Est. No.

STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN
CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

| | |
|---|--|
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice. |
| In case of emergency: Call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor. | |
| Note To Physician: Treat the patient symptomatically. Symptoms of Poisoning. The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by breathing difficulties and sedation. | |

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Socks plus footwear

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

ACCEPTED
with COMMENTS
In EPA Letter Dated
JUN 23 2010
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
264-735

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Do not contaminate feed or food. Keep out of reach of children. Obtain prompt medical aid if poisoning should occur.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

GROUND WATER ADVISORY: Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface, and where the soils are very permeable, i.e., well drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)

Non-refillable Containers

Non-refillable containers - Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Refillable Containers

Refillable container – Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows. Refill this container with pesticide only. Do not reuse this container for any other purpose. After emptying product from container, either return container to Bayer CropScience per instructions from Bayer CropScience Customer Service Center (1-800-527-4781) or rinse and either recycle or dispose of the container as follows:

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs).

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To triple rinse the containers before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

Refillable Containers

End users are authorized to remove tamper evident cables as required to remove the product from the container unless the container is equipped with one way valves and refilling or returning is planned. If this is the case, end users are not authorized to remove tamper evident cables, one way valves or clean container. See Container Disposal instructions under Storage and Disposal.

GENERAL INFORMATION

Sencor® 4 cannot be applied with low pressure or high volume hand wand equipment.

MIXING

When using Sencor® 4, make sure the sprayer is completely clean, free of rust or corrosion which occurs from Winter storage. Examine strainers and screens to be sure the sprayer is clean from previously used pesticides.

Any tank mix containing Sencor® 4 should be kept agitated and sprayed out immediately. Do not allow tank mixes to stand for prolonged periods of time.

The proper mixing procedure for Sencor® 4 alone or in tank mix combinations with other herbicides is:

1. Fill the spray tank 1/4 to 1/3 full with clean water.
2. Add specified rate of Sencor® 4 while recirculating and with agitator running.
3. Follow the triple rinse procedure described under STORAGE AND DISPOSAL to insure that all product is removed from the container.
4. Mix thoroughly and add clean water to fill spray tank to desired level.
5. Add the other herbicide to tank last and agitate thoroughly.
6. Continue agitation during application and until sprayer tank is empty.

SOIL TEXTURE: As used on this label, "Coarse soils" are loamy sand or sandy loam soils. "Medium soils" are loam, silt loam, silt, sandy clay, or sandy clay loam. "Fine soils" are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

GENERAL PRECAUTIONS AND RESTRICTIONS

- Do not use on other crops grown for food or forage. Apply this product only as specified on this label. Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.
- Do not allow sprays to drift on to adjacent desirable plants.
- Observe all cautions and limitations on labeling of all products used in mixtures.
- For all uses: Low pressure and high volume hand wand equipment is prohibited.

CHEMIGATION

Sencor® 4 is recommended for application through sprinkler irrigation equipment to potatoes, soybeans, tomatoes, and asparagus as directed on this label. Refer to the crop sections of this label for rates, weeds controlled or suppressed, restrictions, and special precautions.

Apply this product only through sprinkler (including center pivot, lateral move, or solid set) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

Calibration: (Center Pivot and Self-Propelled Lateral Move Systems): Sprinkler irrigation systems must be accurately calibrated for application of Sencor® 4. Greater accuracy in calibration (and distribution) will be achieved by injecting a larger volume of a more dilute mixture of product and water per hour. Follow the steps below to calibrate center pivot and lateral move systems:

1. Determine number of minutes required to make one complete revolution while applying 1/4 to 3/4 inch of water per acre.
2. With the system at operating pressure determine the exact number of minutes required to inject one gallon of water.
3. Divide the time required for one revolution (step 1) by the time required to inject one gallon (step 2). This gives total gallons of product-water mixture to be added to nurse tank.
4. Add required amount of water to nurse tank and start the agitation system. Then add sufficient Sencor® 4 at the specified rate (See *BROADCAST APPLICATION INSTRUCTIONS*) to the nurse tank.

EXAMPLE: If 20 hours (1200 minutes) were required for one revolution and if 2 minutes were required to inject one gallon, then a total of 600 gallons of product-water mixture are required (1200 / 2 = 600); to treat 135 acres at 1 pint/acre, 135 pints (16 gallons and 7 pints) of Sencor® 4 are required.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in the injection nurse tanks during the herbicide application, sufficient to keep herbicide in suspension.

Apply specified dosage in 1/4 to 3/4 inch of water (1/4 to 1/2 inch of water on sandy soils) per acre as a continuous injection in center pivot and lateral move systems or in the last 15 to 30 minutes of set in permanent solid set sprinkler systems. Application of more than the quantity of irrigation water recommended on this label may result in decreased product performance by removing the chemical from the zone of effectiveness. Where sprinkler distribution patterns do not overlap sufficiently unacceptable weed control may result.

Where sprinkler distribution patterns overlap excessively crop injury may result. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. To insure that lines are flushed and free of remaining pesticide, an indicator dye may be injected into the lines to mark the end of the application period.

Use a minimum of 1 part water to 1 part herbicide for injection. The use of a larger volume of water will insure greater accuracy and more uniform distribution.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
3. Where states have more stringent regulations, they should be observed.
4. The applicator should be familiar with and take into account the information covered in the *Aerial Drift Reduction Advisory Information*.

INFORMATION ON DROPLET SIZE: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see *Wind, Temperature and Humidity, and Temperature Inversions*).

CONTROLLING DROPLET SIZE:

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.


SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

WIND: Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS: Sencor® 4 should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).



APPLICATION OF SENCOR® 4 WITH HERBICIDE SPRAY EQUIPMENT

Use a standard low pressure (20 to 40 psi.) herbicide boom sprayer equipped with suitable nozzles and screens no finer than 50-mesh in-nozzle and in-line strainers. Agitate thoroughly before and during application with bypass agitation.

GROUND APPLICATION: Apply the proper rate of Sencor® 4 in a minimum of 10 to 40 gallons of spray mixture per acre broadcast.

Banded Application: Use proportionally less Sencor® 4 per acre in a band versus a broadcast application. For band application use 1/4 to 1 gallon of spray mix per inch of band width regardless of row spacing.

EXAMPLES: (1) To treat a 15-inch band on rows 30 inches apart, use one-half of the broadcast rate of Sencor® 4. (2) To treat a 14-inch band on rows 42 inches apart, use one-third of the broadcast rate of Sencor® 4.

AERIAL APPLICATION: Where permitted, apply specified rate in a minimum of 2 to 10 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

NOTE: Do not apply aerially when Sencor® 4 is tank mixed with Lasso®.

For All Applications of Sencor® 4: Sprayer must be accurately calibrated before applying Sencor® 4. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases. Avoid over application, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result). Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops, such as cole crops, onions, peas, or strawberries are present in adjacent fields or in areas where wheat is growing in coarse textured soils.

SPRAYER CLEANUP: Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution of Sencor® 4 from the spray tank and dispose of according to label disposal instructions. Rinse the spray tank and refill with water, adding a heavy-duty detergent at the rate of one cup per 20 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles or spray rig. All rinse water must be disposed of in compliance with local, state, and Federal guidelines.

APPLICATION OF SENCOR® 4 IN FLUID FERTILIZERS

Sencor® 4 may be applied in fluid fertilizer solutions to alfalfa and soybeans by following the appropriate mixing procedures and compatibility check. When using tank mix combinations, be sure all components are compatible.

Compatibility checks of Sencor® 4 and tank mix combinations which include Sencor® 4 should be made for each batch of fluid fertilizer because of the variability of these fertilizers.

Compatibility Check:

1. Pre-mix 2 teaspoonfuls of Sencor® 4 with 8 teaspoonfuls of water (1:4 ratio) in a quart jar by adding the water first and followed with Sencor® 4. Mix thoroughly. If a second herbicide is to be used, double the amount of water (1:8 ratio) and add the second herbicide after mixing Sencor® 4 first.
2. Then pour 1 pint of fluid fertilizer into the quart jar and shake well.
3. Allow to stand for 5 minutes.

THIS COMPATIBILITY CHECK SHOULD ONLY BE USED WHEN MIXING WITH FLUID FERTILIZERS.

Interpretation of Results: If the solution in the jar appears to be uniform, without signs of agglomeration, or without a separation of an oily film on top of the fertilizer, the mixture may be used. If not, repeat the compatibility check using twice the amount of water or add a compatibility agent to the water. If separation occurs, but the mixture can be resuspended by shaking, then application is possible with good agitation in the spray tank.

Tank Mixing Guidelines:

1. Add the required amount of water and compatibility agent (if required) to the tank. Start agitation system while adding Sencor® 4 and follow by adding the fluid fertilizer and agitate.
2. If a second herbicide is to be used, follow as above in 1, but use twice the amount of water. Start agitation and add Sencor® and follow by adding the second herbicide, and then continue filling the tank with fluid fertilizer.
3. Maintain continuous agitation to assure uniform spray mixture until the tank is emptied.

COMMERCIAL IMPREGNATION AND APPLICATION OF SENCOR® 4 ON DRY BULK FERTILIZER

Dry bulk fertilizer may be impregnated or coated with Sencor® 4 for application to established alfalfa and to soybeans. All instructions, cautions, and special precautions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling.

Impregnation: To impregnate, use a system consisting of a belt, conveyor, or closed drum which is used for dry bulk fertilizer blending. Any commonly used fertilizer can be impregnated with Sencor® 4 except ammonium nitrate, or fertilizers containing ammonium nitrate, potassium nitrate, or sodium nitrate. Do not use on powder limestone.

Apply using a minimum of 200 lbs dry bulk fertilizer per acre and up to a maximum of 450 lbs per acre. To impregnate or coat dry bulk fertilizer, mix Sencor® 4 with sufficient water to form a sprayable slurry. The delivery nozzles must be directed to deliver a fine spray toward the fertilizer for thorough coverage while avoiding spray contact with mixing equipment. Uniform impregnation of Sencor® 4 to dry bulk fertilizer will vary and if the absorptivity is not adequate, an absorbent powder may be added to produce a dry, free-flowing mixture. Micro-Cel E (Johns-Manville Product Corporation) is the recommended absorbent powder. When another herbicide is used with Sencor® 4, mix and impregnate immediately.

Apply immediately after impregnation unless experience has shown that impregnated fertilizer can be stored without becoming lumpy and difficult to spread.

Rates: Select the specified rate of Sencor® 4 per acre from the appropriate section of this label and refer to the formula below to determine the amount of Sencor® which is to be impregnated on a ton of dry bulk fertilizer based on the amount of fertilizer which will be distributed on one acre.

$$\frac{\text{Pt Sencor® 4}}{\text{Per Acre}} \times \frac{2000 \text{ lb Fertilizer}}{\text{Acre}} = \frac{\text{Pt Sencor® 4}}{\text{Ton of Fertilizer}}$$

APPLICATION: Uniform application is essential for satisfactory weed control. Accurate calibration of fertilizer application equipment is essential for uniform distribution to the soil surface. The recommended method of application is to apply 1/2 the specified rate and overlap 50 percent or to double apply by splitting the middles to obtain the best distribution pattern.

If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

INCORPORATION AND COMBINATION USES: When Sencor® 4 is to be used in combination with another herbicide, follow directions on this label for combinations, rates, crops, incorporation, and special precautions.

SOYBEANS (Except California)

Sencor® herbicide tank mix combinations are recommended for preplant incorporated applications, preemergence surface applications, split-shot application and extended split-shot application. Sencor® 4 is also recommended as an overlay application following a preplant incorporated application of a recommended grass herbicide and alone as a preemergence surface application. All these applications can be applied with ground equipment, and some can be applied with aerial spray equipment. In addition, Sencor® 4 can be applied as a post emergence directed spray to soybeans in certain states.

SPECIAL PRECAUTIONS (Soybeans): Injury to soybeans may occur when Sencor® 4 is used under the following conditions:

- When soils have a calcareous surface area or a pH of 7.5 or higher.
- Due to the sensitivity of certain soybean varieties, do not use Sencor® on Altona, AP 55, AP 71, Asgrow 6520, Burlison, Coker 102, Coker 156, Dassel, GL 3202, Govan, Maple Amber, NB 3665, NKS 1884, Paloma 350, Portage, Regal, Semmes, Terra-Vig 505, Terra-Vig 606, Tracy, Vansoy, and Vinton 81. Consult your Bayer CropScience representative or your seed supplier for information on the tolerance of newly released soybean varieties, prior to use of Sencor®.
- When applied in conjunction with soil-applied organic phosphate pesticides.
- Over application or boom overlapping may result in stand loss and soil residues.
- Uneven application or improper incorporation can decrease the level of weed control and/or increase the level of injury.
- When applied to any soil with less than 1/2% organic matter.
- Soil incorporation deeper than recommended.
- When sprayers are not calibrated accurately.
- When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
- When soybeans are planted less than 1 1/2 inches deep, particularly in preemergence application.


Activation: A minimum amount of soil moisture is required to activate Sencor®. In areas of low rainfall, preemergence applications to dry soil should be followed with light irrigation of 1/4 acre inch of water. Do not apply heavy irrigation immediately after application. As with many surface applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

Grazing and Feeding Treated Vines: Treated vines may be grazed or fed to livestock 40 days after application when Sencor® is applied alone or with Treflan®, Prowl® or Lasso®.

Do not use treated vines for feed or forage when Sencor® is applied with Sonalan®, linuron plus Lasso.

Rate Ranges: Where a rate range is shown, use a lower rate on soils that are coarse textured or low in organic matter. Use a higher rate on soils that are relatively fine textured or high in organic matter.

Replanting: If replanting is necessary in fields treated with Sencor® 4 as directed on this label, the field may be replanted to soybeans. When replanting a minimum of tillage is recommended. Do not apply a second treatment as injury to soybeans may occur.



WEEDS CONTROLLED BY SENCOR® AND SENCOR® HERBICIDE TANK-MIX COMBINATIONS

C = Control

S = Suppression or Erratic Control

P = Poor or No Control

0 = No Information (Control may range from poor to excellent)

1 = Sencor® Alone

4 = Sencor® plus Prowl

7 = Sencor® plus Sonalan

2 = Sencor® Split-Shot

5 = Sencor® plus Lasso

8 = Sencor® plus linuron plus Lasso

3 = Sencor® plus Treflan

6 = Extended Split-Shot

| ANNUAL BROADLEAF WEEDS | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|--|---|---|---|---|---|---|---|---|
| Black Nightshade (<i>Solanum nigrum</i>) | P | P | P | P | C | C | P | S |
| Bristly Starbur (<i>Acanthospermum hispidum</i>) | C | C | C | C | C | C | C | C |
| Buffalobur (<i>Solanum rostratum</i>) | C | C | P | P | P | C | P | O |
| Carpetweed (<i>Mollugo verticillata</i>) | C | C | C | C | C | C | C | C |
| Cocklebur (<i>Xanthium pensylvanicum</i>) | S | C | S | S | S | C | S | S |
| Copperleaf, Hophornbeam (<i>Acalypha ostryaefolia</i>) | C | C | C | C | C | C | C | C |
| Florida Beggarweed (<i>Desmodium tortuosum</i>) | C | C | C | C | C | C | C | C |
| Florida Pusley (<i>Richardia scabra</i>) | C | C | C | C | C | C | C | C |
| Galinsoga (<i>Galinsoga</i> spp.) | C | C | C | C | C | C | C | C |
| Horseweed, Maretail (<i>Conyza canadensis</i>) | O | O | O | O | O | C | O | O |
| Jimsonweed (<i>Datura stramonium</i>) | C | C | C | C | C | C | C | S |
| Knotweed (<i>Polygonum</i> spp.) | C | C | C | C | C | C | C | C |
| Kochia (<i>Polygonum</i> spp.) | C | C | C | C | C | C | C | C |
| Lambsquarters (<i>Chenopodium</i> spp.) | C | C | C | C | C | C | C | C |
| Morningglory, Ivyleaf (<i>Ipomoea hederacea</i>) | P | P | S | P | P | P | P | P |
| Morningglory, Pitted (<i>Ipomoea lacunosa</i>) | P | P | S | P | P | P | P | P |
| Morningglory, Smallflower (<i>Jacquemontia tamnifolia</i>) | P | P | C | P | P | P | P | P |
| Morningglory, Tall (<i>Ipomoea purpurea</i>) | P | P | S | P | P | P | P | P |
| Pigweeds (<i>Amaranthus</i> spp.) | C | C | C | C | C | C | C | C |
| Prickly Sida/ Teaweed (<i>Sida spinosa</i>) | C | C | C | C | C | C | C | C |
| Purslane (<i>Portulaca oleracea</i>) | C | C | C | C | C | C | C | C |
| Ragweed, Common (<i>Ambrosia artemisiifolia</i>) | C | C | C | C | C | C | C | C |
| Redweed (<i>Melochia corchorifolia</i>) | C | C | C | C | C | C | C | C |
| Russian Thistle (<i>Salsola kali</i>) | C | C | C | C | C | C | C | C |
| Sesbania (<i>Sesbania</i> spp.) | C | C | C | C | C | C | C | C |
| Shepherdspurse (<i>Capsella bursa-pastoris</i>) | C | C | C | C | C | C | C | C |
| Sicklepod (<i>Cassia obtusifolia</i>) | C | C | S | S | C | C | S | S |
| Smartweeds (<i>Polygonum</i> spp.) | C | C | C | C | C | C | C | S |
| Spotted Spurge (<i>Euphorbia maculata</i>) | C | C | P | P | C | C | P | O |
| Spurred Anoda (<i>Anoda cristata</i>) | C | C | C | C | C | C | C | O |
| Sunflower (<i>Helianthus</i> spp.) | C | C | S | S | S | C | S | P |
| Velvetleaf (<i>Abutilon theophrasti</i>) | C | C | C | C | C | C | C | C |
| Venice Mallow (<i>Hibiscus trionum</i>) | C | C | C | C | C | C | C | C |
| Wild Mustards (<i>Brassica</i> spp.) | C | C | C | C | C | C | C | C |

| PINTS OF SENCOR® 4 PER ACRE | | | |
|---|-------------------------|----------------|----------------|
| SOIL TEXTURE | ORGANIC MATTER | | |
| | Less than 2% | 2 to 4% | Over 4% |
| COARSE SOILS (Sandy loam, loamy sand) | DO NOT USE ³ | 3/4 | 1 |
| MEDIUM SOILS¹ (Loam, silt loam, silt, sandy clay, sandy clay loam) | 3/4 to 1 | 1 to 1-1/4 | 1-1/4 to 1-1/2 |
| FINE SOILS¹ (Silty clay, silty clay loam ² , clay, clay loam) | 1 to 1-1/4 | 1-1/4 to 1-1/2 | 1-1/2 to 1-3/4 |
| Mississippi Delta Only | 1-1/2 | 1-3/4 | 2 |

¹ For control of lambsquarters, redroot pigweed and wild mustard, and for suppression of green, yellow and giant foxtails on alkaline (calcareous) soils in Nebraska, Minnesota, South Dakota and North Dakota only, apply Sencor® 4 at rates of 1/2 pt/acre on medium soils and 1/2 to 3/4 pt/acre on fine soils regardless of soil organic matter percentage (use 3/4 pt only where soil pH is less than 7.5 and weed pressure is heavy). The 1/2 pt/acre rate of Sencor® 4 alone can be applied regardless of soil pH. For control of other weeds listed on this label use Sencor® 4 at full rates specified in the table above, **but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.**

² Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

³ Refer to the appropriate section of this label for use of Sencor® 4 on soybeans in coarse soils with 0.5% or more organic matter in certain states.

USES OF SENCOR® HERBICIDE IN COMBINATION WITH OTHER HERBICIDES

SEQUENTIAL APPLICATION OF SCEPTER FOLLOWING SENCOR® 4

If needed, application of Sencor® alone or in a registered tank mix according to directions on this label, may be followed by an early postemergence application of Scepter herbicide (1.5 lb/gal liquid or 70 DG) for control of cocklebur. Apply 1/6 to 1/3 pint of Scepter (0.7 to 1.4 ounces of Scepter 70 DG) in a minimum of 20 gallons of water per acre. Use 1/6 pint of Scepter (0.7 ounce of Scepter 70 DG) if cockleburs are less than 3 inches tall or have fewer than 3 leaves and are actively growing. For cockleburs less than 6 inches tall and actively growing use 1/3 pint of Scepter (1.4 ounces of Scepter 70 DG) per acre. Do not use Scepter when soybeans or cockleburs have been subjected to stress conditions such as temperature or moisture extremes. Do not exceed a total of 2/3 pint of Scepter (2.8 ounces of Scepter 70 DG) per acre in one season. Wait at least 10 days after application of Scepter before cultivating.

When preparing the spray mixture with Scepter, add 2 pints of nonionic surfactant approved for use on growing crops and containing at least 80% active ingredient per 100 gallons of mixture. Apply crop oil concentrate (COC) at the rate specified on the COC label.

Use Scepter only in the states where it is registered as listed on the product label.

Apply Scepter at least 90 days before harvest of soybeans. Do not graze or feed soybean forage, hay, or straw to livestock.

Refer to the Scepter label for additional cautions and precautions, directions, limitations, and information on environmental hazards and planting of rotational crops.

SPLIT-SHOT APPLICATION

A preplant incorporated application of Sencor® 4 tank mixed with either Treflan, Lasso, Prowl or Sonalan and followed by a preemergence surface application of Sencor® alone after planting but prior to soybean emergence, will control more broadleaf and grass weeds in soybeans than when either herbicide is used alone.

Refer to the Treflan, Lasso, Prowl or Sonalan labels, and to appropriate sections of this label for directions on soil preparation, herbicide application, incorporation techniques, herbicide rates, weed species controlled, and restrictions for using tank mix combinations of Sencor®. Carefully observe the *Special Precautions* section concerning the use of Sencor® in tank mix combinations on soybeans.

When a Split-Shot application of Sencor® 4 with Prowl, Treflan, or Sonalan is used, the preplant incorporated tank mix may be applied up to 21 days prior to planting soybeans; with Lasso, the preplant incorporated tank mix may be applied up to 14 days prior to planting.

On medium and fine textured soils with greater than 2% organic matter, a rate range is specified for the Sencor® 4 preemergence overlay application. The higher rate should be used (a) in fields with a history of severe broadleaf weed pressure, (b) when the time between preplant incorporated tank mix and preemergence overlay applications approaches the maximum stated above, and/or (c) when the organic matter content of the soil is at the upper end of the indicated range.

For black nightshade control, refer to the appropriate sections of the Lasso or Sonalan labels for specific instructions.

SPLIT-SHOT APPLICATION

Preplant Incorporated Tank-Mix Application - FOLLOWED BY - Preemergence Overlay Application

| SOIL TEXTURE ¹ | Rate of Combination Product/Acre | Plus | Rate of Sencor® 4 Pt/Acre | Rate of Sencor® 4 Pt/Acre ORGANIC MATTER | | |
|--|---|------|--|--|------------------------------|---|
| | | | | Less than 2% | 2 to 4% | Over 4% |
| COARSE (Light) sand, loamy sand, sandy loam | Treflan 1 pt OR Lasso 2 to 2-1/2 qt OR Prowl 1-1/2 pt OR Sonalan 1-1/4 to 2 pt | plus | 1/2-FOLLOWED BY | 1/4 | 1/4 | 1/4 to 1/2 |
| MEDIUM loam, silt loam, sandy clay loam, silt, sandy clay | Treflan 1-1/2 pt OR Lasso 2-1/2 to 3 qt OR Prowl 1-1/2 pt OR Sonalan 1-3/4 to 2-1/2 pt | plus | 3/4-FOLLOWED BY or 1/2 ² -FOLLOWED BY | 1/4 1/2 | 1/4 to 1/2 1/2 to 3/4 | 1/2 to 3/4 (3/4 to 1) ³ |
| FINE (Heavy) silty clay loam*, clay loam, silty clay, clay | Treflan 2 pt OR Lasso 2-1/2 to 3 qt OR Prowl 1-1/2 to 2 pt OR Sonalan 2-1/4 to 3 pt | plus | 1.0-FOLLOWED BY or 3/4 ² -FOLLOWED BY | 1/4 1/2 | 1/4 to 1/2 1/2 to 3/4 | 1/2 to 3/4 (3/4 to 1) ³ |

- * Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.
- ¹ On **coarse textured** soils, do not use on sand soils with less than 1% organic matter, or on loamy sand or sandy loam soils with less than 0.5% organic matter. However, on coarse textured soils **with a calcareous surface area or a pH of 7.5 or higher**, do not use on sand soils with less than 2% organic matter, or on loamy sand or sandy loam soils with less than 1% organic matter.
- ² Use this lower rate of Sencor® 4 in the preplant incorporated tank mix **on soils having a calcareous surface area or a pH of 7.5 or higher**, and in those situations where soils within a field vary extremely in texture or organic matter content.
- ³ Reduce this preemergence overlay rate of Sencor® 4 by 1/4 pt/acre when using SPLIT-SHOT application **on soils with over 4% organic matter and which have a calcareous surface area or a pH of 7.5 or higher**.

EXTENDED SPLIT-SHOT APPLICATION
(Includes No-Till, Reduced-Till, Ridge-Till, Strip-Till, Mulch-Till)

An early preplant (surface-applied or shallow incorporated) application of Sencor® tank mixed with Lasso, followed by a preemergence surface application of Sencor® tank mixed with Lasso after planting but prior to soybean emergence, will control more broadleaf and grass weeds in soybeans than either herbicide used alone.


An Extended Split-Shot application will decrease the need for tillage and/or contact herbicides for the control of existing vegetation prior to planting, while providing residual control of weeds after planting.

When an Extended Split-Shot application of Sencor® 4 with Lasso is used, the preplant tank mix combination may be applied 15 to 30 days prior to planting soybeans. Follow directions on the label accompanying the product for Split-Shot applications from 0 to 14 days before planting.

Where a rate range is specified, the higher rates should be used (a) in fields with a history of severe weed pressure, (b) when the time between early preplant tank mix and preemergence overlay applications approaches the maximum 30 days, (c) when the organic matter content of the soil is at the upper end of the indicated range, (d) when heavy crop residues are present on the soil surface, and/or (e) when the early preplant tank mix application is shallow incorporated (e.g. use 2 to 2 1/2 qt Lasso in the early preplant tank mix when surface applied and use 2 1/2 to 3 qt Lasso when the tank mix is to be lightly incorporated).

When weeds exceed 1 to 1 1/2 inches in height or diameter at application, use a contact herbicide, such as Round-Up® or Gramoxone.

Refer to the Lasso label, and to appropriate sections of this label for additional information on soil preparation, herbicide application, weeds controlled, precautions, restrictions, limitations and sprayer clean up.



| EXTENDED SPLIT-SHOT APPLICATION | | | | | | | | | |
|--|----------------------------------|------|---------------------------|-------------|----------------------------------|------|--|------------|----------|
| Early Preplant Tank Mix Application (Surface-Applied or Shallow Incorporated) | | | | Followed By | Preemergence Overlay Application | | | | |
| SOIL TEXTURE ¹ | Rate of Combination Product/Acre | Plus | Rate of Sencor® 4 Pt/Acre | | Rate of Combination Product/Acre | Plus | Rate of Sencor® 4 Pt/Acre ORGANIC MATTER | | |
| | | | | | | | 1/2 to 2% | 2 to 4% | Over 4% |
| COARSE (Light) (Sand, loamy sand, sandy loam) | Lasso 1-1/2 to 2 qt | plus | 1/2 to 3/4 | Lasso | 2/3 pt 1-1/2 qt | plus | 1/4 | 1/4 to 1/2 | 1/2 |
| MEDIUM (Loam, silt loam, sandy clay loam, silt, sandy clay) | Lasso 2 to 3 qt | plus | ² 3/4 to 1 | Lasso | 3/4 pt 1 to 2 qt | plus | 1/2 | 1/2 to 3/4 | 3/4 to 1 |
| FINE (Heavy) (Silty clay loam*, clay loam, silty clay, clay) | Lasso 2 to 3 qt | plus | ² 1 to 1-1/4 | Lasso | 1 pt 1 to 2 qt | plus | 1/2 | 1/2 to 3/4 | 3/4 to 1 |

* Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

¹ On **coarse textured** soils, do not use on sand soil with less than 1% organic matter. However, on coarse textured soils **with a calcareous surface area or a pH of 7.5 or higher**, do not use on sand soils with less than 2% organic matter, or on loamy sand or sandy loam soils with less than 1% organic matter.

² Use the lower rate of Sencor® 4 in the early preplant tank mix **on soils having a calcareous surface area or a pH of 7.5 or higher, and in those rare situations where soils within a field vary extremely in texture or organic matter content.**

SENCOR® PLUS SONALAN

Sencor® plus Sonalan Overlay Application: Sencor® 4 may be applied as a preemergence overlay application following a preplant incorporated application of Sonalan 3 EC. Consult the Sonalan label for specific directions on use, instructions, restrictions and any additional weeds not specified on this label.

Sencor® plus Sonalan Tank Mix Application: Incorporate the tank mixture into the top 1 to 2 inches of soil within 21 days before planting according to label directions for Sonalan.

Apply Sencor® 4 plus Sonalan preplant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If soybeans are planted on beds, apply and incorporate the tank mixture after bed formation.

Mixing: Refer to the *General Information* section in the front of this label.

Application: Sonalan should be uniformly applied and thoroughly mixed into the soil within 2 days after application. For specific application information, refer to the *General Information* section in the front of this label.

| BROADCAST RATES | | |
|--|-------------------|---------------------|
| SOIL TEXTURE | Sencor® 4 Pt/Acre | Sonalan 3EC Pt/Acre |
| COARSE ¹ (Sandy loam, loamy sand) | 1/2 | 1 1/4 to 2 |
| MEDIUM ³ (Loam, silt loam, silt, sandy clay, sandy clay loam) | 3/4 | 1 3/4 to 2 1/2 |
| FINE ³ (Silty clay, silty clay loam ² , clay, clay loam) | 1 | 2 1/4 to 3 |

¹ Do not use on coarse soils with less than 1% organic matter.

² Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

³ For control of lambsquarters, redroot pigweed, wild mustard, and green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Sencor® 4 at rates of 1/2 pt/acre on medium soils and 1/2 to 3/4 pt/acre on fine soils regardless of soil organic matter percentage (use 3/4 pt only where soil pH is less than 7.5 and weed pressure is heavy). The 1/2 pt rate of Sencor® 4 in tank mix combination with Sonalan can be applied regardless of soil pH. For control of other weeds not listed on the label, use Sencor® 4 at full rates specified in the table above, **but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.**

SPECIAL PRECAUTIONS (Sencor® plus Sonalan):

- For additional precautions, restrictions, limitations, incorporation, and sprayer clean-up information, refer to the appropriate sections of this label and the Sonalan label.
- For black nightshade control, refer to the Sonalan label for specific rates and application instructions.

SENCOR® PLUS TREFLAN

Sencor® and Treflan Overlay Application: Sencor® 4 may be applied as a preemergence broadcast or band overlay application following a pre-plant incorporated treatment of Treflan. Consult the Treflan label for specific directions for use, instructions, restrictions and any additional weeds not specified on this label.

Sencor® plus Treflan Tank Mix Application: A single application of a tank mix combination of Sencor® 4 and Treflan EC will control more broadleaf and grass weeds in soybeans than when either herbicide is used alone.

Prepare the soil surface by deep plowing, offset disking or tandem disking prior to the application of the herbicide combination. The soil surface should be well prepared and free of clods and trash.

This Sencor® plus Treflan tank mix combination may be applied and incorporated into the soil up to 10 days before planting.

Mixing: Refer to the *General Information* section in the front of this label.

Application: For specific application information refer to the *General Information* section in the front of this label.

Apply Sencor® 4 plus Treflan to the soil surface and incorporate in the same operation, if possible. Variable weed control may result from delayed incorporation if Sencor® plus Treflan are applied to a wet, warm soil surface or if the wind velocity is 10 miles per hour or higher. Use machinery that mixes Sencor® plus Treflan thoroughly with the soil. Incorporation may be delayed up to 24 hours after application. Shallow incorporation with implements set to cut less than 2 inches deep may result in erratic weed control. Do not use spike or spring-tooth harrows alone for incorporation.

| BROADCAST RATES | | |
|--|-----------------------|------------------------|
| SOIL TEXTURE | Sencor® 4 Pt Per Acre | Treflan EC Pt Per Acre |
| COARSE ¹ (Sandy loam, loamy sand) | 1/2 | 1 |
| MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | 3/4 | 1-1/2 |
| FINE (Silty clay, silty clay loam ² , clay, clay loam) ³ | 1 | 2 |

¹ Do not use on coarse soils with less than 1% organic matter.

² Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

³ For control of lambsquarters, redroot pigweed, wild mustard, and green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Sencor® 4 at rates of 1/2 pt/acre on medium soils and 1/2 to 3/4 pt/acre on fine soils regardless of soil organic matter percentage (use 3/4 pt only where soil pH is less than 7.5 and weed pressure is heavy). The 1/2 pt rate of Sencor® 4 in tank mix combination with Treflan can be applied regardless of soil pH. For control of other weeds listed on the label use Sencor® 4 at full rates specified in the table above, **but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.**

Incorporation Equipment:

Set PTO driven equipment (tillers, cultivators, hoes) to cut 2 to 3 inches deep and space rotors to provide a clean sweep of the soil. PTO equipment should not be operated at a speed greater than 4 miles per hour.

Set disk to cut 4 to 6 inches deep and operate twice in different directions at 4 to 6 miles per hour.

Set mulch treader and other similar disk-type implements to cut 3 to 4 inches deep and operate twice in different directions at 5 to 8 miles per hour.

For Coarse and Medium Textured Soils Only:

Set rolling cultivator to cut 2 to 4 inches deep and operate twice at 6 to 8 miles per hour. Set bed conditioner (Do-All) to cut 2 to 4 inches deep and operate at 4 to 6 miles per hour.

SPECIAL PRECAUTIONS (Sencor® plus Treflan):

- Seedling disease, cold weather, excessive moisture, high salt concentration or drought may weaken soybean seedlings and increase possibility of damage from the tank mix. Do not plant soybeans deeper than 2 inches.
- Do not rotate any crop not listed on this label for 18 months after the tank-mix application.
- For additional precautions, restrictions, limitations and sprayer clean-up information refer to the appropriate section of this label. Do not use this tank mix combination on soils containing charcoal in Arkansas, Louisiana and Mississippi.

SENCOR® PLUS PROWL

Sencor® plus Prowl Overlay Application: Apply a preplant incorporated treatment of Prowl as directed on that product label for use on soybeans. Follow with a preemergence treatment of Sencor® 4 as directed on this label for use on soybeans.

Sencor® plus Prowl Tank Mix Application

Preplant Incorporated Application: Prepare the soil by plowing or disking to mix previous crop residues into the soil to a depth of 4 to 6 inches.

For specific application information refer to the "General Information" section in the front of this label.

Incorporate the tank mixture into the top 1 or 2 inches of soil within 7 days after application according to label directions for Prowl. Mechanical incorporation is not required if a rain of one-quarter inch or more occurs within 7 days after application. Soybeans must be planted no later than 7 days after application of the tank mixture.

Preemergence Application: Except for minimum and no-tillage systems, the seed bed should be firm and free of trash and clods.

For specific application information refer to the "General Information" section in the front of this label. Do not apply Prowl preemergence north of Interstate 80.

This application must be made after planting and before crop emergence. Do not incorporate.

If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tilling equipment such as a rotary hoe that does not damage soybeans.

Mixing Instructions: Refer to the "General Information" section in the front of this label.

For information on applying Sencor® 4 in fluid or dry fertilizer refer to the "Application Of Sencor® 4 In Fluid Fertilizers" or "Commercial Impregnation And Application Of Sencor® 4 On Dry Bulk Fertilizer" under the "General Information" section in the front of this label.

SOUTHERN STATES AND EASTERN COASTAL PLAINS

For use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, Southeastern Missouri "Bootheel" Region and Coastal Plains of Delaware*, Maryland*, New Jersey* and Virginia*.

* Do not use Sencor®/Prowl on soils with less than 2% organic matter in the coastal plain of New Jersey or the Delmarva Peninsula.


BROADCAST RATES
Sencor® Plus Prowl Tank Mix Applications

| SOIL TEXTURE | Sencor® 4 Pt/Acre | Prowl Pt/Acre |
|--|-------------------|---------------|
| COARSE ¹ (Sandy loam, loamy sand) | 1/2 | 1-1/2 |
| MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | 3/4 | 1-1/2 |
| FINE (Silty clay, silty clay loam ² , clay, clay loam) | 1 | 1-1/2 to 2 |

¹ Do not use on sand soils. Do not use on loamy sand or sandy loam containing less than 1% organic matter.

² Silty clay loam soils are transitional soils and may be classified as medium textured soils in certain regions of the U.S.

Do not use on muck or peat soils.



NORTHEASTERN AND NORTH CENTRAL STATES

For use only in Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New York, North Dakota, Ohio, Pennsylvania, South Dakota, Wisconsin and Missouri (except the "Bootheel" Region).

BROADCAST RATES
Sencor® Plus Prowl Tank Mix Applications

1/2 to 3% ORGANIC MATTER

| SOIL TEXTURE | Sencor® 4 Pt/Acre | Prowl Pt/Acre |
|--|-------------------|---------------|
| COARSE ¹ (Sandy loam, loamy sand) | 1/2 | 1 |
| MEDIUM (Loam, silt loam, sandy clay, sandy clay loam) | 3/4 | 1-1/2 to 2 |
| FINE (Silty clay, silty clay loam ² , clay, clay loam) | 3/4 to 1 | 1-1/2 to 2 |

Over 3% ORGANIC MATTER

| | | |
|--|------------|------------|
| COARSE ¹ (Sandy loam, loamy sand) | 3/4 | 1-1/2 |
| MEDIUM (Loam, silt loam, sandy clay, sandy clay loam) | 3/4 to 1 | 1-1/2 to 2 |
| FINE (Silty clay, silty clay loam ² , clay, clay loam) | 1 to 1-1/4 | 2 to 2-1/2 |


¹ Do not use on sand soils. Do not use on loamy sand or sandy loam containing less than 1% organic matter. Where a range of rates is shown for medium and fine soils, use the higher rate if heavy weed infestations are anticipated.

² Silty clay loam soils are transitional soils and may be classified as medium textured soils in certain areas of the U.S.

Do not use on muck or peat soils.

SPECIAL PRECAUTIONS (Sencor® plus Prowl):

- Soil incorporation deeper than recommended will reduce weed control and can result in crop injury.
- For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the Prowl label.



SENCOR® PLUS LASSO

Sencor® Plus Lasso Tank Mix Application:

PREEMERGENCE

Sencor® 4 may be used in a tank mix combination with Lasso as a preemergence band or broadcast application to soybeans in accordance with the specified soil types and dosages.

For specific information regarding spray equipment, dilution rates, mixing, directions for use, methods of application, limitations and restrictions refer to the appropriate section of this label.


Refer to the Lasso label for pertinent instructions, directions for use, restrictions and any additional weeds not specified on this label.

| APPLICATION INSTRUCTIONS | | | |
|---|-------------------|------|---------------|
| Sencor® Plus Lasso Tank Mix Preemergence Application (Broadcast Rates) | | | |
| 1/2 to 3% Organic Matter | | | |
| SOIL TEXTURE | Sencor® 4 Pt/Acre | Plus | Lasso Qt/Acre |
| COARSE ¹ (Sandy loam) | 1/2 | plus | 1-1/2 to 2 |
| MEDIUM ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 3/4 | plus | 1-1/2 to 2 |
| FINE ² (Silty clay, silty clay loam ³ , clay, clay loam) | 1 | plus | 2 |
| MISSISSIPPI DELTA ONLY (Silty clay to heavy clay) | 2 | plus | 2 to 2-1/2 |
| Greater than 3% Organic Matter | | | |
| COARSE ¹ (Sandy loam) | 3/4 | plus | 1-1/2 to 2 |
| MEDIUM ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1 | plus | 1-1/2 to 2 |
| FINE ² (Silty clay, silty clay loam ³ , clay, clay loam) | 1 to 1-1/4 | plus | 2 to 2-1/2 |
| MISSISSIPPI DELTA ONLY (Silty clay to heavy clay) | 2 | plus | 2 to 2-1/2 |

¹ Do not use Sencor® plus Lasso on sand or loamy sand soils with less than 2% organic matter.

² For control of lambsquarters, redroot pigweed, wild mustard, green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Sencor® 4 at rates of 1/2 pt/acre on medium soils and 1/2 to 3/4 pt/acre on fine soils regardless of soil organic matter percentage (use 3/4 pt only where soil pH is less than 7.5 and weed pressure is heavy). The 1/2 pt/acre rate of Sencor® 4 in tank mix combination with Lasso can be applied regardless of soil pH. For control of other weeds use Sencor® 4 at full rates specified in the table above, **but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.**

³ Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.



PREPLANT INCORPORATED: For specific application information refer to the "General Information" section in the front of this label. Apply Sencor® 4 plus Lasso preplant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If soybeans are planted on beds, apply and incorporate the tank mixture after bed formation. Apply within 7 days prior to planting and shallowly incorporate into the upper 1 to 2 inches of soil.

| APPLICATION INSTRUCTIONS | | |
|---|-------------------|---------------|
| Sencor® Plus Lasso Tank Mix Preplant Incorporated Applications (Broadcast Rates) | | |
| SOIL TEXTURE | Sencor® 4 Pt/Acre | Lasso Qt/Acre |
| COARSE ¹ (Loamy sand [over 2% organic matter], sandy loam) | 1/2 | 2 to 2-1/2 |
| MEDIUM (Loam, silt loam, silt) | 3/4 | 2-1/2 to 3 |
| FINE (Silty clay loam ² , sandy clay loam, silty clay, sandy clay, clay loam, clay) | 1 | 2-1/2 to 3 |
| MISSISSIPPI DELTA ONLY (Silty clay, clay) | 1 to 1-1/4 | 2-1/2 to 3 |

¹ Do not use Sencor® plus Lasso on sand or loamy sand soils with less than 2% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.
Do not use on muck soils.

SPECIAL PRECAUTIONS (Sencor® Plus Lasso):

- For additional precautions, restrictions, limitations and sprayer clean up information, refer to the appropriate sections of this label and the Lasso label.

SENCOR® PLUS COMMAND®

Sencor® 4 may be applied in combination with Command 4EC as a preplant or shallow incorporated application for the control of certain weeds in soybeans. Consult the Command label for specific directions on use, instructions, restrictions and any additional weeds not specified on this label.

Mixing: Refer to the "General Information" section in the front of this label.

Application: Sencor® plus Command may only be applied with ground equipment as a preplant or shallow incorporated application. Sencor® plus Command should be immediately incorporated into the top 1 to 3 inches after application unless surface is dry. On dry soils, incorporate into the top 1 to 3 inches within 3 hours of tank mix application.

Do not apply this tank mix within 1000 feet of towns and subdivisions, commercial vegetable, fruit, nurseries or greenhouse operations.


A minimum of 15 gallons spray volume per acre is recommended with appropriate nozzle types and sizes to produce a coarse spray droplet. The use of an approved agricultural drift reducing additive is recommended for application volumes of 15 to 40 gallons per acre. The use of an approved agricultural drift reducing additive is required at spray volumes of 10 to 15 gallons per acre.

NOTE: Off-site movement of Command spray drift or vapors can cause foliar whitening or yellowing of some vegetation. Prior to application of Command, read and strictly follow all precautions and application instructions as set forth in that label.

For additional information on application, refer to the "General Information" section in the front of this label and the Command label.

Weeds Controlled: Bristly Starbur, Carpetweed, Copperleaf, Florida Beggarweed, Florida Pusley, Galinsoga, Jimsonweed, Knotweed, Lambsquarters, Pigweeds, Prickly Sida/Teaweed, Purslane, Common Ragweed, Redweed, Sesbania, Smartweeds, Spurred Anoda, Velvetleaf, Venice Mallow, Wild Mustards, Barnyardgrass*, Bluegrass, Broadleaf Signalgrass, Crabgrass*, Foxtails (Green, Giant, Yellow*, Robust Purple), Goosegrass, Johnsongrass (seedling)*, Fall Panicum*, Texas Panicum, and witchgrass.

- * Use 2 pt/A Command 4EC on coarse and medium textured soils with high populations of these weeds.



| APPLICATION INSTRUCTIONS | | |
|---|---------------------|---------------------|
| Sencor® 4 Plus COMMAND 4EC Tank Mix Preplant Incorporated Application (Broadcast Rates) | | |
| SOIL TEXTURE ¹ | Sencor® 4 (Pt/Acre) | COMMAND 4EC Pt/Acre |
| 0.5% to 3% ORGANIC MATTER | | |
| COARSE² (Sandy loam, loamy sand) | 1/2 | 1-1/2 to 2 |
| MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 to 3/4 | 1-1/2 to 2 |
| FINE (Silty clay, silty clay loam ³ , clay, clay loam) | 1/2 to 3/4 | 1-1/2 to 2 |
| Over 3% ORGANIC MATTER | | |
| COARSE² (Sandy loam, loamy sand) | 1/2 | 1-1/2 to 2 |
| MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 to 3/4 | 1-1/2 to 2 |
| FINE (Silty clay, silty clay loam ³ , clay, clay loam) | 3/4 to 1 | 1-1/2 to 2 |

¹ Crop injury may occur on soils having a calcareous surface area or a pH of 7.1 or higher.
² Do not use on coarse soils with less than 1% organic matter.
³ Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

SPECIAL PRECAUTIONS (Sencor® 4 plus Command):

- Do not rotate to wheat, barley, alfalfa or seed corn in the fall of the year of application or in the spring of the following year as crop injury may occur. Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.
- Do not apply when weather conditions favor drift. Do not use treated vines for feed or forage.
- Do not apply aerially or through irrigation equipment.

SENCOR® PLUS COMMENCE®

Sencor® plus Commence Tank Mix Early Preplant Incorporated Application: Sencor® 4 in a tank mix with Commence 5.25 EC may be applied broadcast preplant incorporated up to 30 days before planting soybeans for the control of certain broadleaf weeds and grasses. Refer to the Commence herbicide label for additional directions for use, weeds controlled, instructions, restrictions and limitations not specified on this label.

Mixing: Refer to the "General Information" section on this label.

Application: For information on applying Sencor® 4, refer to the "General Information" section on this label.

| APPLICATION INSTRUCTIONS | | |
|---|-------------------|--------------------------|
| Sencor® 4 Plus Commence 5.25 EC Tank Mix Early Preplant Incorporated Application ¹ | | |
| SOIL TEXTURE | Sencor® 4 Pt/Acre | Commence 5.25 EC Pt/Acre |
| 1/2 to 3% ORGANIC MATTER | | |
| COARSE² | 3/4 | 1-1/3 to 2 |
| MEDIUM | 1 | 2 to 2-1/4 |
| FINE | 1 | 2-2/3 |
| Over 3% ORGANIC MATTER | | |
| COARSE² | 3/4 | 1-1/3 to 2 |
| MEDIUM | 1 | 2 to 2-1/4 |
| FINE | 1-1/2 | 2-2/3 |

¹ For use on soils with a pH of 7.5 or lower.
² Do not use on coarse soils with less than 1% organic matter.

RESTRICTIONS AND LIMITATIONS:

- Do not apply aerially or through irrigation equipment.
- Do not apply when weather conditions favor drift. Do not allow sprays to drift on to adjacent desirable plants. Do not use treated vines for feed or forage.
- Do not rotate to wheat, barley, alfalfa or seed corn in the fall of the year of application or in the spring of the following year as crop injury may occur. Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.

Sencor® plus Commence Tank Mix Preplant Incorporated: Sencor® 4 may be tank mixed with Commence 5.25 EC for preplant incorporated application to control certain weeds in soybeans. Refer to the "General Information" section of this label for information on mixing, application, restrictions, special precautions and weeds controlled by Sencor® 4. See appropriate sections of the Commence 5.25 EC herbicide label for additional precautionary statements, directions for use, instructions and additional weeds controlled.


| APPLICATION INSTRUCTIONS | | |
|---|---------------------|----------------------------|
| Sencor® Plus Commence Tank Mix Preplant Incorporated Application (Broadcast Rates) | | |
| SOIL TEXTURE | Sencor® 4 (Pt/Acre) | Commence 5.25 EC (Pt/Acre) |
| 1/2 to 3% ORGANIC MATTER | | |
| COARSE ² (Sandy loam, loamy sand) | 1/2 | 1-1/3 to 2 |
| MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 to 3/4 | 2 to 2-1/4 |
| FINE (Silty clay, silty clay loam ³ , clay, clay loam) | 1/2 to 3/4 | 2-2/3 |
| Over 3% ORGANIC MATTER | | |
| COARSE ² (Sandy loam, loamy sand) | 1/2 | 1-1/3 to 2 |
| MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 to 3/4 | 2 to 2-1/4 |
| FINE (Silty clay, silty clay loam ³ , clay, clay loam) | 3/4 to 1 | 2-2/3 |
| ¹ Crop injury may occur on soils having a calcareous surface area or a pH of 7.1 or higher. ² Do not use on coarse soils with less than 1% organic matter. ³ Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S. | | |

SPECIAL PRECAUTIONS (Sencor® plus Commence):

- Do not rotate to wheat, barley, alfalfa or seed corn in the fall of the year of application or in the spring of the following year as crop injury may occur. Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.
- Do not apply when weather conditions favor drift. Do not use treated vines for feed or forage. Do not apply aerially or through irrigation equipment.
- Do not allow sprays to drift on to adjacent desirable plants.

SENCOR® 4 PLUS FREEDOM®

Sencor® 4 may be tank mixed with Freedom 3 EC for preplant incorporated application to control weeds in soybeans. Refer to the "General Information" section of this label for information on mixing, application, restrictions, special precautions and weeds controlled by Sencor® 4. See appropriate sections of the Freedom 3 EC herbicide label for additional precautionary statements, directions for use, instructions and additional weeds controlled.



APPLICATION INSTRUCTIONS

Sencor® plus Freedom Tank Mix Preplant Incorporated Application (Broadcast Rates)

| SOIL TEXTURE | Sencor® 4 Lb/Acre | Plus | Freedom 3 EC Qt/Acre |
|---|----------------------|------|-------------------------|
| 1/2 to 3% Organic Matter | | | |
| COARSE ¹ (Sandy loam) | 1/2 | Plus | 2-3/4 to 3-1/2 |
| MEDIUM ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 3/4 | Plus | 2-3/4 to 3-1/2 |
| FINE ² (Silty clay, silty clay loam ³ , clay, clay loam) | 1 | Plus | 3-1/2 to 4 |
| MISSISSIPPI DELTA ONLY (Silty clay to heavy clay) | 2 | Plus | 3-1/2 to 4-1/2 |
| Greater than 3% Organic Matter | | | |
| COARSE ¹ (Sandy loam) | 3/4 | Plus | 3 to 3-1/2 |
| MEDIUM ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1 | Plus | 3-1/2 to 4 |
| FINE ² (Silty clay, silty clay loam ³ , clay, clay loam) | 1 to 1-1/4 | Plus | 3-1/2 to 4-1/2 |
| MISSISSIPPI DELTA ONLY (Silty clay to heavy clay) | 2 | Plus | 3-1/2 to 4-1/2 |

¹ Do not use Sencor® plus Freedom on sand or loamy sand soils with less than 2% organic matter.

² For control of lambsquarters, redroot pigweed, wild mustard, green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Sencor® 4 at rates of 1/2 pt/acre on medium soils and 1/2 to 3/4 pt/acre on fine soils regardless of soil organic matter percentage (use 3/4 pt only where soil pH is less than 7.5 and weed pressure is heavy). The 1/2 pt/acre rate of Sencor® 4 in tank mix combination with Freedom can be applied regardless of soil pH. For control of other weeds use Sencor® 4 at full rates specified in the table above, **but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.**

³ Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

SPECIAL PRECAUTIONS (Sencor® plus Freedom):

- Do not use on muck soils.
- Do not allow sprays to drift on to adjacent desirable plants.

SENCOR® PLUS CANOPY® PLUS A GRASS HERBICIDE

A tank mix combination of Sencor® 4 plus Canopy 75 DF plus a registered and recommended grass herbicide (Lasso, Prowl, Sonalan or Treflan) is recommended for control of the following weeds in soybeans:

WEEDS CONTROLLED

| Annual Grasses | | | |
|-------------------------|---------------|-------------------------|----------------|
| Barnyardgrass | Crabgrass | Johnsongrass (seedling) | Sandbur |
| Bluegrass | Crowfootgrass | Junglerice | Sprangletop |
| Broadleaf signalgrass | Foxtails | Panicum, Fall | Stinkgrass |
| Browntop Millet | Goosegrass | Panicum, Texas | |
| Annual Broadleaves | | | |
| Bristly Starbur | Galinsoga | Prickly Sida/Teaweed | Shepherdspurse |
| Carpetweed | Jimsonweed | Purslane | Smartweed |
| Cocklebur | Knotweed | Ragweed, Common | Spurred Anoda |
| Copperleaf, Hophornbeam | Kochia | Redweed | Velvetleaf |
| Florida Beggarweed | Lambsquarters | Russian Thistle | Venice mallow |
| Florida Pusley | Pigweed | Sesbania | Wild mustard |

Tank mix combinations which include Lasso or Prowl can be applied preemergence broadcast or preplant incorporated broadcast. When Sonalan or Treflan are used in the tank mix, apply preplant incorporated broadcast. Refer to the table below for rates of each product to be used in tank mix combinations:

| APPLICATION INSTRUCTIONS | | | |
|--|---------------------------|-------------------------|-------------------------|
| Sencor® 4 Plus Canopy 75 DF Plus A Grass Herbicide (Broadcast Rates) | | | |
| Product | SOIL TEXTURE ¹ | | |
| | COARSE ² | MEDIUM | FINE |
| Sencor® 4 (Pt/Acre) | 1/2 | 1/2 to 3/4 ³ | 3/4 to 1/3 ³ |
| Canopy DF (Oz/Acre) | 3 | 3 | 3 to 4 |
| Treflan (Pt/Acre) | 1 | 1-1/2 | 2 |
| Prowl (Pt/Acre) | 1-1/2 | 1-1/2 to 2 | 1-1/2 to 2-1/2 |
| Lasso (Qt/Acre) | 2 to 2-1/2 | 2-1/2 to 3 | 2-1/2 to 3 |
| Sonalan (Pt/Acre) | 1-1/4 to 2 | 1-1/4 to 2-1/2 | 2-1/4 to 3 |

¹ Do not use on soils with a pH greater than 7.0.
² Refer to "Soil Texture" paragraph on this label for specific soil classification.
³ Use the lower rate of Sencor® 4 in preplant incorporated tank-mix as in those situations where soils within a field vary extremely in texture or organic matter content.

Refer to the "General Information" section of this label for mixing and application directions.

IMPORTANT: If weeds escape in fields treated with these tank mix combinations, postemergence application of a registered and recommended herbicide will be needed for control.

SPECIAL PRECAUTIONS:

- For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate sections of the labels for Sencor® 4 and Canopy 75 DF.
- Do not use treated vines or feed or forage.

SENCOR® PLUS COMMAND PLUS A GRASS HERBICIDE

Sencor® 4 may be applied with Command 4EC and a grass herbicide (Treflan, Lasso, Prowl or Sonalan) for the control of certain broadleaf weeds and grasses in soybeans. This combination will provide improved control of heavy infestations of velvetleaf, jimsonweed and common ragweed. Sencor® 4 and Command 4EC plus a grass herbicide may be applied preplant incorporated broadcast. Consult the Command, Treflan, Lasso, Prowl or Sonalan labels for specific directions for use, instructions, restrictions and additional weeds controlled not specified on this label.

Mixing: Refer to the "General Information" section in the front of this label.

Application: For specific application information, refer to the "General Information" section in the front of this label.

WEEDS CONTROLLED

| Annual Grasses | | | |
|-------------------------|----------------------|-------------------------|----------------|
| Barnyardgrass | Browntop millet | Foxtails | Panicum, Fall |
| Bluegrass | Crabgrass | Goosegrass | Witchgrass |
| Broadleaf signalgrass | Crowfootgrass | Johnsongrass (seedling) | |
| Annual Broadleaves | | | |
| Bristly Starbur | Knotweed | Ragweed, Common | Smartweeds |
| Carpetweed | Kochia | Redweed | Spotted spurge |
| Copperleaf, Hophornbeam | Lambsquarters | Russian Thistle | Spurred Anoda |
| Florida Beggarweed | Pigweeds | Sesbania | Velvetleaf |
| Florida Pusley | Prickly Sida/Teaweed | Shepherdspurse | Venice mallow |
| Galinsoga | Purslane | Sicklepod | Wild Mustards |
| Jimsonweed | | | |

Sencor® and Command plus Treflan, Lasso, Prowl or Sonalan will provide suppression (reduce the competition) of cocklebur and sunflower.

| APPLICATION INSTRUCTIONS | | | |
|---|---------------------------|-------------------------|-----------------------|
| Sencor® 4 plus Command plus a Grass Herbicide (Broadcast Rates) | | | |
| Product | SOIL TEXTURE ¹ | | |
| | COARSE | MEDIUM | FINE |
| Sencor® 4 (Pt/Acre) | 1/2 | 1/2 to 3/4 ² | 3/4 to 1 ² |
| Command 4EC ³ (Pt/Acre) | 1/2 to 3/4 | 1/2 to 3/4 | 1/2 to 3/4 |
| Treflan (Pt/Acre) | 1 | 1-1/2 | 2 |
| Prowl (Pt/Acre) | 1-1/2 | 1-1/2 to 2 | 1-1/2 to 2-1/2 |
| Lasso (Qt/Acre) | 2 to 2-1/2 | 2-1/2 to 3 | 2-1/2 to 3 |
| Sonalan (Pt/Acre) | 1-1/4 to 2 | 1-3/4 to 2-1/2 | 2-1/4 to 3 |

¹ Refer to "Soil Texture" paragraph on this label for specific soil classification. On coarse textured soils **with a calcareous surface area or a pH of 7.5 or higher**, do not use on loamy sand or sandy loam soils with less than 1% organic matter.

² The higher rate of Sencor® 4 is recommended for the control of sicklepod and hemp sesbania. Use the lower rate of Sencor® 4 in the preplant incorporated tank mix **on soils having a calcareous surface area or a pH of 7.5 or higher**, and in those situations where soils within a field vary extremely in texture or organic matter content.

³ Higher rate is recommended under moderate to heavy weed infestations.

SENCOR® PLUS SCEPTER PLUS A GRASS HERBICIDE

Sencor® 4 may be applied with Scepter herbicide and a grass herbicide (Treflan, Lasso, Prowl or Sonalan) for the control of certain broadleaf weeds and grasses in soybeans. Sencor® and Scepter plus Treflan or Sonalan may be applied preplant incorporated broadcast. Sencor® and Scepter plus Lasso or Prowl may be applied preplant incorporated, preemergence broadcast or in a band application.

Consult the Scepter, Treflan, Lasso, Prowl or Sonalan labels for specific directions for use, instructions, restrictions and additional weeds controlled not specified on this label.

Mixing: Refer to the "General Information" section in the front of this label.


Application: For specific application information, refer to the "General Information" section in the front of this label.

Weeds Controlled: Sencor® 4 plus Scepter plus Treflan, Lasso, Prowl or Sonalan will control the following broadleaf weeds and grasses:

WEEDS CONTROLLED

| Annual Grasses | | | |
|-------------------------|---------------------------|-------------------------|----------------|
| Barnyardgrass | Browntop Millet | Foxtails | Panicum, Fall |
| Bluegrass | Crabgrass | Goosegrass | Witchgrass |
| Broadleaf signalgrass | Crowfootgrass | Johnsongrass (seedling) | |
| Annual Broadleaves | | | |
| Bristly Starbur | Galinsoga | Prickly Sida/Teaweed | Smartweeds |
| Buffalobur | Jimsonweed | Purslane | Spotted spurge |
| Carpetweed | Knotweed | Ragweed, Common | Spurred Anoda |
| Cocklebur | Kochia | Redweed | Sunflower |
| Coffee Senna | Lambsquarters | Russian Thistle | Velvetleaf |
| Copperleaf, Hophornbeam | Morningglory, pitted | Sesbania | Venice mallow |
| Florida Beggarweed | Morningglory, smallflower | Shepherdspurge | Wild mustards |
| Florida Pusley | Pigweeds | Sicklepod | |

Sencor® and Scepter plus Treflan, Lasso, Prowl or Sonalan will suppress (reduce the competition of) ivyleaf and tall morningglory, and red rice.



| Sencor® 4 plus Scepter plus a Grass Herbicide (Broadcast Rates) | | | |
|---|---------------------------|-------------------------|-----------------------|
| Product | SOIL TEXTURE ¹ | | |
| | COARSE | MEDIUM | FINE |
| Sencor® 4 (Pt/A) | 1/2 | 1/2 to 3/4 ² | 3/4 to 1 ² |
| Scepter (1.5 lb/Gal liquid ³ Pt/A) - or - | 1/3 to 1/2 | 1/3 to 1/2 | 1/3 to 1/2 |
| Scepter 70 DG ³ (Oz/A) | 1.4 to 2.1 | 1.4 to 2.1 | 1.4 to 2.1 |
| Treflan (Pt/A) | 1 | 1-1/2 | 2 |
| Prowl (Pt/A) | 1-1/2 | 1-1/2 to 2 | 1-1/2 to 2-1/2 |
| Lasso (Qt/A) | 2 to 2-1/2 | 2-1/2 to 3 | 2-1/2 to 3 |
| Sonalan (Pt/A) | 1-1/4 to 2 | 1-3/4 to 2-1/2 | 2-1/4 to 3 |

¹ Refer to "Soil Texture" paragraph on this label for specific soil classification. On coarse textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on loamy sand or sandy loam soils with less than 1% organic matter.

² The higher rate of Sencor® is recommended for preemergence tank mix application and for the control of sicklepod and hemp sesbania. Use the lower rate of Sencor® in the preplant incorporated tank mix on soils having a calcareous surface area or a pH of 7.5 or higher, and in those situations where soils within a field vary extremely in texture or organic matter content.

³ Higher rate is recommended under moderate to heavy weed infestations.

SENCOR® PLUS PURSUIT® AND A GRASS HERBICIDE

Sencor® 4 may be tank mixed with Pursuit herbicide and a registered and recommended grass herbicide (Lasso, Prowl, Sonalan or Treflan) for control of certain broadleaf and grass weeds in soybeans. Refer to the product labels for Pursuit, Lasso, Prowl, Sonalan or Treflan for additional directions for use, instructions, restrictions and limitations not included on this label.

Tank mix combinations of Sencor® 4, Pursuit and Lasso or Prowl can be applied broadcast preemergence or preplant incorporated. When the grass herbicide used is Sonalan or Treflan, apply the tank mix broadcast preplant incorporated.

Mixing and Application: Refer to the "General Information" section of this label for directions on mixing and application of Sencor® 4.

| APPLICATION INSTRUCTIONS | | |
|---|---------------------|-------------------|
| Sencor® 4 Plus Pursuit and a Grass Herbicide* | | |
| Soil Texture | Sencor® 4 (Pt/Acre) | Pursuit (Oz/Acre) |
| COARSE | 1/2 | 4 |
| MEDIUM | 3/5 to 3/4 | 4 |
| FINE | 3/4 to 1 | 4 |


*For control of grass weeds, include Lasso, Prowl, Sonalan or Treflan at label rates in the tank mix with Sencor® 4 and Pursuit herbicides.

Restrictions and Limitations: Do not apply this tank mix with aerial or irrigation equipment. Do not apply when weather conditions favor drift, or allow sprays to drift on to adjacent desirable plants. Do not use treated vines for feed or forage. Refer to appropriate sections of the Pursuit herbicide label for restrictions on use area and rotational crops.

SENCOR® PLUS PURSUIT PLUS HERBICIDE

Sencor® 4 may be tank mixed with Pursuit Plus herbicide for broadcast preemergence or preplant incorporated application to soybeans for control of certain broadleaf and grass weeds. Refer to the Pursuit Plus herbicide label for additional directions for use, instructions, restrictions and limitations not included on this label.

Mixing and Application: Refer to the "General Information" section of this label for directions on mixing and application of Sencor® 4.



| APPLICATION INSTRUCTIONS | | |
|--|--------------------|-----------------------|
| Sencor® 4 plus Pursuit Plus Herbicide (Broadcast Rates) | | |
| Soil Texture | Sencor® 4 (P/Acre) | Pursuit Plus (P/Acre) |
| COARSE | 1/2 | 2-1/2 |
| MEDIUM | 3/5 to 3/4 | 2-1/2 |
| FINE | 3/4 to 1 | 2-1/2 |

Restrictions and Limitations:

- Do not apply this tank mix with aerial or irrigation equipment. Do not apply when weather conditions favor drift, or allow sprays to drift on to desirable plants.
- Do not use treated vines for feed or forage.
- Refer to appropriate sections of the Pursuit Plus herbicide label for restrictions on use area and rotational crops.

SENCOR® PLUS LINURON PLUS LASSO

Sencor® plus linuron plus Lasso Tank Mix Application: Sencor® 4 may be applied in combination with linuron 50 DF or 4L and Lasso 4 as a preemergence application for the control of certain weeds in soybeans. Consult the linuron or Lasso labels for specific directions for use, instructions, restrictions and any additional weeds not specified on this label.

Mixing: Refer to the "General Information" section in the front of this label.

Application: Applications can be made only with ground spray equipment in accordance with specified soil types and dosage rates. For specific application information, refer to the "General Information" section in the front of this label.

| Sencor® Plus Linuron Plus Lasso Broadcast Rates (0.5 to 3% Organic Matter Only) | | | |
|--|---|---|---|
| Product | SOIL TEXTURE | | |
| | COARSE ¹ (Sandy, loamy sand, sandy loam) | MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam) | FINE (Silty clay, silty clay loam ² , clay, clay loam) |
| Sencor® 4 (pt/acre) | 1/4 to 3/8 | 3/8 to 1/2 | 1/2 to 3/4 |
| Linuron 50 DF (lb/acre) or Linuron 4L (pt/acre) | 1/3 to 1/2 | 1/2 to 3/4 | 3/4 to 1-1/2 |
| Lasso 4 (qt/acre) | 3/4 to 1 | 1 to 1-1/2 | 1-1/4 to 2 |

¹ Do not use Sencor® plus linuron plus Lasso on sand soils with less than 1% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.


SPECIAL PRECAUTIONS (Sencor® plus Linuron plus Lasso: For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the linuron or Lasso labels.

FOR USE IN COARSE (LIGHT) SOILS IN Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia.

Sencor® 4 herbicide is recommended alone or in combination with Treflan or Lasso for use in coarse-textured, low organic matter soils in the states listed above for the control of certain weeds in soybeans. Refer to the appropriate sections of this label and the Treflan or Lasso labels for specific directions for use, instructions, restrictions and any additional weeds not specified on this label.

Mixing: Refer to the "General Information" section in the front of this label.

Application: For specific application information, refer to the "General Information" section in the front of this label.



Sencor® (Alone) Preemergence Application (Broadcast Rates)

| SOIL TEXTURE | ORGANIC MATTER | Sencor® 4 Pt/ACRE |
|--|----------------|-------------------------|
| COARSE (LIGHT) SOILS Sand ¹ , Loamy Sand, Sandy Loam | 0.5% or Above | 1/2 to 3/4 ² |

¹ Do not use on sand with less than 1% organic matter.
² Use the higher rate under heavy weed pressures and/or on soils higher in organic matter.

Sencor® in Combination with Other Herbicides: Sencor® 4 is recommended in a tank mix combination with Treflan as a preplant incorporated application or as a preemergence overlay application following a preplant incorporated application of Treflan. Sencor® 4 is also recommended for use as a preemergence application in combination with Lasso.

**FOR USE IN COARSE (LIGHT) SOILS
0.5% or Above ORGANIC MATTER
(Broadcast Rates)**

| SOIL TEXTURE | COMBINATION PRODUCT/ACRE | Plus | Sencor® 4 Pt/ACRE |
|---|--|------|-------------------------|
| COARSE (LIGHT) SOILS Sand ¹ , Loamy sand, Sandy loam | Preplant Incorporated Treflan 4EC 1 pt | Plus | 1/2 to 3/4 ² |
| | Preemergence Lasso 4E 1-1/2 to 2 qt | Plus | 1/2 to 3/4 ² |

¹ Do not use on sand with less than 1% organic matter.
² Use the higher rate under heavy weed pressures and/or on soils higher in organic matter.

SPECIAL PRECAUTIONS:

- Do not use on sand soils with less than 1% organic matter, or on sandy loam or loamy sand soils with less than 0.5% organic matter.
- For additional precautions, restrictions, limitations and sprayer clean-up information, refer to the appropriate sections of this label and the Treflan or Lasso labels.

BURNDOWN WEED CONTROL - FIELD CORN AND SOYBEANS

Sencor® 4 can be used as part of a herbicide program for burndown of existing vegetation prior to crop emergence in conservation tillage systems. Sencor® may be tank mixed with 2,4-D low volatile ester (LVE), Gramoxone Extra, or Roundup / Roundup Ultra / Touchdown for control of emerged weeds prior to field corn or soybean emergence.

Sencor® tank mixes with 2,4-DB, Fusion, Poast Plus or Select may also be used in soybeans for control of emerged weeds prior to crop emergence. Sencor® 4 burndown tank mixes can be applied before planting or prior to crop emergence in the following areas:


Field Corn:

Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota, and Wisconsin

Soybeans:

All areas for all products except Fusion tank mixes - see Fusion section of this label for specified states.

Application: Sencor® 4 may be applied up to 30 days prior to planting or preemergence. Apply only by ground equipment when Sencor® 4 is used for burndown of existing vegetation in conservation tillage systems. Sencor® 4 and tank mix partner burndown rates are listed in the following three tables.



| SENCOR® 4 BURNDOWN RATES - FIELD CORN AND SOYBEANS | | |
|---|---------------------------|----------------------|
| CROPS | APPLICATION TIMING | SENCOR® 4 RATE (P/A) |
| Field corn Iowa Kansas Missouri Nebraska South Dakota | Preplant (0 to 30 days) | 3 to 8 |
| | Preemergence | |
| Field corn Illinois Indiana Kentucky Michigan Minnesota Ohio Wisconsin | Preplant (10 to 30 days) | 3 to 8 |
| | Preplant (0 to 9 days) | 3 to 6 |
| | Preemergence | |
| Soybeans | Preplant (0 to 30 days) | 3 to 8 |
| | Preemergence | |

| SENCOR® 4 PLUS TANK MIX PARTNER BURNDOWN RATES - FIELD CORN OR SOYBEANS | | |
|--|--|--|
| PRODUCT | RATE | DIRECTIONS AND REMARKS |
| Sencor® 4 + 2,4-D LVE | 3 to 8 fl oz/A* + 1/4 to 1 lb ai/A | In soybeans, apply at least 7 days preplant when using 2,4-D LVE at 1/4 to 1/2 lb ai/A and at least 30 days preplant with rates greater than 1/2 lb ai/A. Include crop oil concentrate (COC) at the rate of 1 gal/100 gal of spray solution (1% v/v). In corn, apply at least 7 days preplant or at least 3 days after planting but before corn emergence. |
| Sencor® 4 + Gramoxone Extra | 3 to 8 fl oz/A* + 24 to 48 fl oz/A | Must be applied prior to crop emergence. Use 24 to 32 fluid ounces of Gramoxone Extra for weeds less than 4 inches in height and 32 to 48 fluid ounces when weeds are 4 to 6 inches in height. Apply in 20 to 60 gallons of water per acre. Include either nonionic surfactant at 1 quart per 100 gallons (0.25% v/v) or crop oil concentrate at 1 gallon per 100 gallons (1% v/v) of spray solution. |
| Sencor® 4 + Gramoxone Extra + 2,4-D LVE | 3 to 8 fl oz/A* + 24 to 48 fl oz/A + 1/4 to 1 lb ai/A | For this tank mix follow the Directions and Remarks Sections above for Sencor® + 2,4-D LVE and Sencor® + Gramoxone Extra, paying special attention to crop planting restrictions with 2,4-D LVE. Include either nonionic surfactant or crop oil concentrate in this tank mix. |
| Sencor® 4 + Roundup/ Roundup Ultra or Touchdown | 3 to 8 fl oz/A* + 12 to 24 fl oz/A or 8 to 16 fl oz/A | Must be applied prior to crop emergence. Use the higher rates as weeds approach the maximum weed heights listed in the "Weeds Controlled" section below. Apply in 10 to 20 gallons of water per acre. With Roundup and Touchdown, include nonionic surfactant at 2 quarts per 100 gallons (0.5% v/v) and ammonium sulfate (spray grade) at 17 pounds per 100 gallons of spray solution. With Roundup Ultra, include ammonium sulfate (spray grade) at 17 pounds per 100 gallons of spray solution. Any glyphosate formulation registered and labeled for use in field corn or soybeans may be tank mixed with Sencor®. |
| Sencor® 4 + Roundup/ Roundup Ultra or Touchdown + 2,4-D LVE | 3 to 8 fl oz/A* + 12 to 24 fl oz/A or 8 to 16 fl oz/A + 1/4 to 1 lb ai/A | For this tank mix follow the Directions and Remarks Sections above for Sencor® + 2,4-D LVE and Sencor® + Roundup / Roundup Ultra / Touchdown, paying special attention to planting restrictions with 2,4-D LVE. Use the adjuvant recommendations under the Sencor® + Roundup / Roundup Ultra / Touchdown tank mix. Do not use crop oil concentrate. |

* If applied to field corn grown in Illinois, Indiana, Kentucky, Michigan, Minnesota, Ohio and Wisconsin, refer to Table 1 for correct Sencor® rate based on application timing.

| SENCOR® 4 PLUS TANK MIX PARTNER BURNDOWN RATES - SOYBEANS ONLY |
|--|
|--|

| PRODUCT | RATE | DIRECTIONS AND REMARKS |
|--|---|--|
| Sencor® 4 + 2,4-DB | 3 to 8 fl oz/A + 1/8 to 7/32 lb ai/A | Apply preplant or before soybean emergence. Include nonionic surfactant at 2 quarts per 100 gallons (0.5% v/v) of spray solution. |
| Sencor® 4 + Fusion + 2,4-D LVE | 3 to 8 fl oz/A + 4 to 8 fl oz/A + 1/4 to 1 lb ai/A | For use only in Delaware, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Pennsylvania, South Dakota, Virginia, West Virginia, and Wisconsin. For this tank mix follow the planting restrictions under the Directions and Remarks Section above for Sencor® + 2,4-D LVE. Fusion rates of 4, 6 and 8 fl. ounces will control certain grasses up to 2, 4 and 6 inches in height, respectively. Include either crop oil concentrate at 1 gallon per 100 gallon (1.0% v/v) or nonionic surfactant at 1 to 2 quarts per 100 gallons (0.25 - 0.5% v/v) of spray solution. Refer to the Fusion label for additional information. |
| Sencor® 4 + Poast Plus + 2,4-D LVE | 3 to 8 fl oz/A + 8 to 12 fl oz/A + 1/4 to 1 lb ai/A | For this tank mix follow the planting restrictions under the Directions and Remarks Section above for Sencor® + 2,4-D LVE. The 8 and 12 fl oz rate of Poast Plus will control certain grasses up to 2 and 3 inches in height, respectively. Include either crop oil concentrate at the rate of 1 gallon per 100 gallons of spray solution (1% v/v) or Dash HC at 1 pint per acre. Refer to the Poast Plus label for additional information. |
| Sencor® 4 + Select + 2,4-D LVE | 3 to 8 fl oz/A + 3 to 4 fl oz/A + 1/4 to 1 lb ai/A | For this tank mix follow the planting restrictions under the Directions and Remarks Section above for Sencor® + 2,4-D LVE. The 3 and 4 fluid ounce rates of Select will control certain grasses up to 3 and 4 inches in height, respectively. Include crop oil concentrate at the rate of 1 quart per acre and 28% UAN (urea ammonium nitrate) at a rate of 1-2 quarts per acre. Refer to the Select label for additional information. |

Special Precautions. Do not apply these treatments after crop emergence. Observe all precautions and limitations on the labeling of all products used in tank mixtures. Refer to the General Information section of this label for additional information, precautions, and limitations.

Field Corn:


- Do not apply on coarse textured soils with less than 1.5% organic matter.
- Do not apply more than 6 fl oz of Sencor® 4 per acre on soils with less than 2% organic matter.
- Do not apply on soils having pH 7.0 or greater.
- Do not apply more than 8 fl oz Sencor® 4 (0.25 pound active ingredient) per acre per growing season.
- Corn seed should be planted a minimum of 1-1/2 inches deep.
- Sencor® 4 may only be used in hybrid seed corn production fields if both inbred parents are known to be tolerant to Sencor® 4.

Soybeans:

- Apply only 2,4-D low volatile ester formulations which are registered and recommended for preplant or burndown use in soybeans.
- Do not apply tank mixtures containing 2,4-D LVE if wind is blowing toward desired susceptible plants (i.e. cotton, tobacco, tomato, etc.) or when wind speeds exceed 6 miles per hour.

Feeding restrictions. Corn treated with Sencor® 4 may be harvested for silage or grain 60 days after treatment. Soybean vines or hay treated with Sencor® 4 may be grazed or fed to livestock 40 days after application. Do not feed hay, forage, fodder or graze 2,4-D, Select, or Fusion treated vegetation. Follow the most restrictive preharvest interval of all products used in a tank mixture.

Weeds controlled: Sencor® 4 in tank mixtures with the above herbicides will provide burndown control of the weeds listed below.



| WEEDS CONTROLLED BY BURNDOWN RATES OF SENCOR® 4 | | | | | | | | | |
|---|----------------------------------|------------------------|--------------------|--------------------|-----------------------------------|---|--------------------|-----------------------------|----------------|
| WEEDS CONTROLLED | SENCOR® 4 PLUS | | | | | | | | |
| | 2,4-D LVE | Poast Plus + 2,4-D LVE | Select + 2,4-D LVE | Fusion + 2,4-D LVE | Roundup/ Roundup Ultra/ Touchdown | Roundup/ Roundup Ultra/ Touchdown + 2,4-D LVE | Gramoxone Extra | Gramoxone Extra + 2,4-D LVE | 2,4-DB |
| BROADLEAVES | MAXIMUM BURNDOWN HEIGHT (INCHES) | | | | | | | | |
| Buffalobur | | | - | | 6 | 6 | 4 to 6 | 4 to 6 | - |
| Chickweed, common | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 2 |
| Cocklebur, common | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 6 |
| Dandelion, common | | | 6 dia ^a | | 2 dia ^b | 6 dia ^a | 4 dia ^d | 6 dia ^a | 2 dia |
| Henbit | | | 4 | | 4 | 4 | 4 to 6 | 4 to 6 | - |
| Horseweed/marestail | | | 6 ^{ac} | | 4 ^b | 6 | 3 | 6 ^a | 2 ^c |
| Jimsonweed | | | 6 | | 6 | 6 | 4 to 6 | 4 to 6 | 2 |
| Kochia* | | | 4 ^{ac} | | 4 | 4 | 4 | 4 | - |
| Ladysthumb | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Lambsquarters, common | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 2 |
| Lettuce, prickly | | | 6 | | 4 | 6 | 4 to 6 | 4 to 6 | 2 |
| Mallow, Venice | | | 6 | | 6 | 6 | 4 to 6 | 4 to 6 | - |
| Morningglory spp. | | | 6 | | 2 | 4 | 2 | 4 | 4 |
| Mustard spp. | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 2 |
| Pennycress, field | | | 6 | | 6 | 6 | 4 to 6 | 4 to 6 | 2 |
| Pigweed, spp. (annual) | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Ragweed, common | | | 6 | | 6 ^b | 8 | 4 to 6 | 4 to 6 | 2 |
| Ragweed, giant | | | 6 ^{ac} | | 4 ^b | 6 | 4 | 6 | 2 |
| Shepherdspurse | | | 6 | | 6 | 6 | 4 to 6 | 4 to 6 | - |
| Sida, prickly | | | 6 | | 4 | 4 | 4 | 4 | 1 |
| Smartweed, Pennsylvania | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Sunflower, common | | | 6 | | 6 | 6 | 4 to 6 | 4 to 6 | 4 |
| Thistle, Russian | | | 4 ^{ac} | | 2 to 4 ^{bc} | 6 | 4 | 4 to 6 | 3 ^c |
| Velvetleaf | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Waterhemp spp. | | | 6 | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |


^a Use 2,4-D LVE at 0.5 pound active ingredient per acre.

^b Use a minimum Roundup/Roundup Ultra rate of 16 fl oz/A and a minimum Touchdown rate of 10.6 fl oz/A.

^c Use Sencor® 4 at 6 fl oz/A for optimum control.

^d Suppression only.

* Does not control triazine resistant biotypes.



| WEEDS CONTROLLED BY BURNDOWN RATES OF SENCOR® 4 (continued) | | | | | | | | | |
|---|----------------------------------|------------------------|--------------------|--------------------|-----------------------------------|---|-----------------|-----------------------------|--------------------------------|
| WEEDS CONTROLLED | SENCOR® 4 PLUS | | | | | | | | |
| | 2,4-D LVE | Poast Plus + 2,4-D LVE | Select + 2,4-D LVE | Fusion + 2,4-D LVE | Roundup/ Roundup Ultra/ Touchdown | Roundup/ Roundup Ultra/ Touchdown + 2,4-D LVE | Gramoxone Extra | Gramoxone Extra + 2,4-D LVE | 2,4-DB |
| ANNUAL GRASSES | MAXIMUM BURNDOWN HEIGHT (INCHES) | | | | | | | | |
| Barley | - | - | - | - | 8 | | 4 to 6 | | Does not control these species |
| Barnyardgrass | 2 to 3 | 3 to 4 | - | - | 6 | | 4 to 6 | | |
| Crabgrass spp. | 2 to 3 | - | - | - | 6 | | 4 to 6 | | |
| Foxtail spp. | 2 to 3 | 3 to 4 | 2 to 6 | - | 8 | | 4 to 6 | | |
| Johnsongrass, seedling | 2 to 3 | - | - | - | 8 | | 4 to 6 | | |
| Panicum, fall | 2 to 3 | 3 | 2 to 6 | - | 6 | | 4 to 6 | | |
| Sandbur, field | - | - | - | - | 8 | | 4 to 6 | | |
| Shattercane | 2 to 3 | - | - | - | 8 | | 4 to 6 | | |
| Wheat, volunteer | - | - | - | - | 6 | | 4 to 6 | | |
| Witchgrass | 2 to 3 | - | - | - | 6 | | 4 to 6 | | |

RESIDUAL WEED CONTROL

Sencor® 4 burndown programs can be used as part of a full season weed control program in both field corn and soybeans when, 1) applied as a tank mixture with residual herbicides, or 2) followed with a postemergence weed control program, which is registered for use on that crop.

For residual control, Sencor® 4 burndown programs may include tank mixes with the following herbicides or combination of herbicides:

Field Corn

| | | | |
|--------------------|--------------|--------------|-----------------|
| Alachlor | Bullet | Harness Xtra | Pursuita |
| Atrazine | Clarity | Lariat | Pursuit Plusa |
| Banvel | Cycle | Linex | Ramrod |
| Bicep | Dual | Linuron | Ramdor/Atrazine |
| Bicep II | Dual II | Lorox | Simazine |
| Bicep Lite | Extrazine II | Marksman | Surpass |
| Bladex | Frontier | Pentagon | Surpass 100 |
| Broadstrike + Dual | Guardman | Prowl | Topnotch |
| Broadstrike Plus | Harness | | |

^a Use only Pursuit resistant/tolerant corn hybrids.

Soybeans

| | | | |
|--------------------|----------|----------------|--------------|
| Alachlor | Dual | New Lorox Plus | Pursuit Plus |
| Broadstrike + Dual | Dual II | Pentagon | Scepter |
| Canopy | Frontier | Preview | Sencorb |
| Command | Gemini | Prowl | Squadron |
| Detail | Linuron | Pursuit | Turbo |

^b Sencor® used (alone and in tank-mixes) on soybeans at higher labeled rates than those listed for burndown weed control will also provide residual control of those weeds listed in the "Weeds Controlled by Sencor® and Sencor® Tank Mix Combinations" section of the Sencor® label.

Refer to the individual product labels for additional information, precautions, and limitations.

SOUTHERN AND SOUTHEASTERN STATES ONLY POSTEMERGENCE DIRECTED SPRAY APPLICATIONS

Sencor® 4 can be applied in postemergence directed sprays to soybeans for control of certain weeds which escape preplant or preemergence herbicide applications and for control of additional flushes of weeds that may occur after soybeans have emerged. Postemergence directed sprays of Sencor® 4 can be applied to soybeans in addition to a preemergence or preplant application of Sencor® herbicide according to label directions.

Weeds Controlled: Sencor® 4, applied postemergence to soybeans as a directed spray according to directions on this label, will control the following at rates shown (broadcast basis) when grasses and common ragweed are less than 1-inch tall and other broadleaves are less than 3-inches tall:

| | | |
|-------------------------|--|---|
| 1/2 Pt/Acre | Florida beggarweed (<i>Desmodium tortuosum</i>) Pigweeds (<i>Amaranthus</i> spp.) Carpetweed (<i>Mollugo verticillata</i>) Cocklebur (<i>Xanthium pensylvanicum</i>) Dayflower (<i>Commelina</i> spp.) | Mexicanweed (<i>Caperonia castaniifolia</i>) Purslane (<i>Portulaca oleracea</i>) Sicklepod (<i>Cassia obtusifolia</i>) Velvetleaf (<i>Abutilon theophrasti</i>) Crabgrass (<i>Digitaria</i> spp.) |
| 1/2 to 1 Pt/Acre | Sesbania (<i>Sesbania</i> spp.) | Prickly sida/Teaweed (<i>Sida spinosa</i>) |
| 1 Pt/Acre | Ragweed, common (<i>Ambrosia artemisiifolia</i>) | |

At the rate of 1/2 pt/acre morningglory species, (*Ipomoea* spp.) horsenettle, (*Solanum* spp.) Florida pusley, (*Richardia scabra*) spotted spurge (*Euphorbia maculata*) and wild poinsettia (*Euphorbia heterophylla*) are suppressed when Sencor® 4 is applied before these weeds are 3 inches tall. The 1 pt/acre rate will suppress broadleaf signalgrass (*Brachiaria platyphylla*) up to 1 inch tall.

SENCOR® POSTEMERGENCE DIRECTED SPRAY

| APPLICATION INSTRUCTIONS | |
|--|-------------------------------|
| CROP | Sencor® 4 Pt/Acre |
| Soybeans (Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee and Texas) | 1/2 to 1 (Broadcast Basis) |

Apply proper dosage using 10 to 40 gallons of water per acre as a directed spray in a 6 to 8 inch band on each side of the row after soybeans are 8 inches tall and before broadleaf weeds are 3 inches tall and before grasses and common ragweed are 1 inch tall. For best results the spray must cover weed foliage with minimum or no contact with soybean foliage. Add a nonionic surfactant such as Ortho X-77 to the spray mixture to obtain better wetting of weed leaf surfaces. To determine the correct dosage of Sencor® 4 for a band application see "Banded Application" under the "General Information" section in the front of this label.

If necessary, a second postemergence directed spray application can be made after 7 days.

Do not feed or graze green soybean vines. Do not harvest soybeans or use dry soybean vines for feed or forage within 70 days of last application.

SPECIAL PRECAUTIONS (Directed Postemergence): Do not apply directly to soybeans or serious crop injury will occur. Do not allow spray to contact more than the lower 1/4 to 1/3 of soybean plants. Soybean leaves contacted by the spray will be killed.

Do not apply Sencor® 4 postemergence to sensitive soybean varieties. See "Special Precautions" in the front of this label.

To avoid injury to other crops or desirable plants from spray drift, sprayer pressure must not exceed 30 psi and the sprayer must be fitted with nozzles no smaller than 8002 T-Jet (or equivalent). Do not apply under weather conditions which favor drift.

POTATOES

Sencor® 4 herbicide is recommended for use in ground, aircraft or specified chemigation equipment as a preemergence and/or postemergence application to potatoes. Early maturing smooth skinned white and all red skinned varieties may be injured with postemergence applications. The varieties Atlantic, Bellchip, Centennial, Chipbelle and Shepody are sensitive to Sencor® 4. Avoid postemergence applications on these varieties. Preemergence applications on these varieties may cause crop injury under adverse weather conditions, on coarse soils, under high soil pH, with higher rates per acre and with mechanical incorporation.

Ground Application: Sencor® 4 is recommended for use with ground spray equipment applied as a preemergence and/or postemergence application for control of the listed grass and broadleaf weeds in potatoes. Apply as a uniform broadcast spray at 20 or more gallons per acre.

Aerial Application: Sencor® 4 may be applied in aerial spray equipment as a preemergence and/or postemergence application at 5 or more gallons per acre.

Chemigation: Sencor® 4 may be applied preemergence and/or early postemergence to potatoes using center pivot, solid set and lateral roll systems. Apply specified dosage in 1/4 to 3/4 inch of water per acre (1/4 to 1/2 inch on sandy soil) as a continuous injection in self-propelled systems or apply in the last 15 to 30 minutes of the set in other systems. Be sure all the Sencor® has been flushed from the lines before shutting down the system.

WEEDS CONTROLLED

Sencor® 4 applied to potatoes according to directions, will provide economic control of the following weeds. For optimum control, applications should be made before weeds are 1 inch tall. (See **NOTE**)

| BROADLEAVES | | | |
|--|----------------------------------|-------------------------------------|--|
| Carpetweed, common ¹ | Mustard, Indian ¹ | Pigweed, redroot ^{1,2} | Sicklepod ¹ |
| Cocklebur, common ^{1,2} | Mustard, tansy ¹ | Pigweed, smooth ^{1,2} | Smartweed, Pennsylvania ^{1,2} |
| Jimsonweed ¹ | Mustard, tumble ¹ | Ragweed, common ^{1,2} | Sunflower, common ³ |
| Kochia ³ | Mustard, wild ¹ | Shepherdspurse ¹ | Thistle, Russian ² |
| Lambsquarters, common ^{1,2} | Pennycress, field ^{1,2} | | |
| GRASSES | | | |
| Barnyardgrass ³ | Foxtail, giant ¹ | Foxtail, yellow ¹ | Panicum, fall ¹ |
| Crabgrass, large ¹ | Foxtail, green ¹ | Johnsongrass, seedling ¹ | Signalgrass, broadleaf ¹ |
| Crabgrass, smooth ¹ | | | |
| ¹ Weeds controlled with preemergence applications. | | | |
| ² Weeds controlled with postemergence applications. | | | |
| ³ Weeds requiring two applications for control. | | | |

HARD TO CONTROL WEEDS

Although Sencor® 4 may not provide commercially acceptable control in every instance, it will suppress growth of the following weeds and reduce their competition with potato plants.

| BROADLEAVES | | | |
|---------------|-------------------|------------------|-------------------|
| Kochia | Nightshade, hairy | Purslane, common | Sunflower, common |
| GRASSES | | | |
| Barnyardgrass | Nutsedge, yellow | | |

NOTE: Where triazine-resistant weeds are present, Sencor® 4 alone may not provide adequate control.

| BROADCAST APPLICATION INSTRUCTIONS | |
|---|-------------------|
| CROP | Sencor® 4 Pt/Acre |
| Potatoes | 1/2 to 2 |
| PREEMERGENCE: Apply specified dosage as a broadcast spray. Do not mechanically incorporate into soil. Use the 1/2 to 1 pint/acre rate for control of wild mustard (<i>Brassica</i> sp.) only. On sand soils or sensitive varieties, do not exceed 1 pint/acre. | |
| Potatoes (Except early maturing smooth skinned, red skinned, and other specified varieties.) | 1/2 to 1 |
| POSTEMERGENCE: Apply specified dosage as a broadcast spray over the tops of potato plants.* Use rates of 1/2 to 1 pint/acre for control of redroot pigweed and common lambsquarters only. Apply the 1 pint/acre rate for control of other weeds listed on this label. | |
| SPLIT APPLICATIONS: This product may be applied once preemergence and once postemergence as directed above.* Do not exceed 2 pint total per acre per season. | |
| IDAHO, OREGON AND WASHINGTON ONLY: Two postemergence applications can be made as broadcast sprays over the tops of potato plants if Sencor® 4 is not applied preemergence. Use 1/2 to 1 pint/acre for control of redroot pigweed and lambsquarters only. On coarse (sandy) soils with low organic matter do not exceed 3/4 pint/acre per application. On medium and heavy soils only, use 1 pint/acre per application for control of other weeds listed on this label and for suppression of hairy nightshade. Make the first application early in the season while weeds are still small. Allow at least 14 days before the second application. Do not apply after June 30 if treated land is to be planted to crops other than potatoes. | |
| * Refer to Special Precautions - Potatoes below. | |

TANK MIXES: Sencor® 4 may be tankmixed with the following herbicides: Dual/Dual II, Eptam, Prowl 3.3 EC and Matrix. In addition, three-way tankmix combinations may be used for Sencor® 4 plus Dual/Dual II, Eptam or Prowl 3.3 EC plus Matrix when applied preemergence. Refer to each product's label for precautionary statements, restrictions, application information and weeds controlled.

Dual/Dual II: Sencor® 4 may be applied in a tankmix combination with Dual/Dual II as a preemergence broadcast application. Apply Sencor® 4 at 3/4 to 2 pints per acre and Dual/Dual II at 1.5 to 3 pints per acre according to the respective labels for use of each product alone on potatoes.

Eptam: Sencor® 4 may be tankmixed with Eptam at rates and uses permitted on each product's label.

Prowl 3.3 EC: Sencor® 4 may be applied in tankmix combination with Prowl as a preemergence or early postemergence broadcast application. As a preemergence mix, apply Sencor® 4 at 1 to 2 pints per acre and Prowl at 1.2 to 3.6 pints per acre. As an early

postemergence spray, apply Sencor® 4 at 1/2 to 1 pint per acre and Prowl at 1.2 to 3.6 pints per acre before the crop is in the 6-inch growth stage.

Matrix: (except the following counties in Colorado: Alamosa, Conejos, Costilla, Rio Grande and Saguache): Sencor® 4 may be applied in tankmix combination with Matrix as a preemergence and/or early postemergence application for improved control on weeds such as Russian thistle, kochia and common lambsquarters. As a preemergence mix, apply Sencor® 4 at 1/2 to 1.125 pints per acre and Matrix at 1 to 1-1/2 oz. product per acre. As an early postemergence spray, apply Sencor® 4 at 1/2 to 1 pint per acre and Matrix at 1 to 1-1/2 oz. product per acre. Use a nonionic surfactant at a rate of 0.125% v/v (1 pt./100 gallon of water). Apply before the crop exceeds 14 inches in height. Postemergence applications of Matrix treatments should be made prior to June 30.

SPECIAL PRECAUTIONS (Potatoes):

- Do not use Sencor® 4 on potatoes in Kern County, California.
- Do not apply more than a total of 2 pints Sencor® 4 per acre in a single crop season regardless of the method of application. Do not make postemergence applications prior to rainfall or irrigation on recently cultivated potatoes, nor within 3 days after periods of cool, wet cloudy weather or injury may occur.
- Postemergence applications may cause some chlorosis or minor necrosis. These symptoms may be more severe if seed-piece decay is occurring or if growing conditions favor crop stress.
- Postemergence applications may be made only on russet or white skinned varieties that are not early maturing.
- Potato varieties may vary in their response to herbicide application. When using Sencor® 4 for the first time on a particular variety, always determine crop tolerance before using on a field scale.
- Do not apply Sencor® 4 within 60 days of harvest.
- Do not use air blast sprayers.
- Do not apply to sweet potatoes or yams.
- Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.

Certain varieties of wheat and barley are sensitive to Sencor® 4 (see the wheat and barley section of this label for sensitive varieties) and should not be planted during the next growing season unless the following cultural practices occur:

1. Potato vines left in rows as a result of harvest must be uniformly distributed over the soil surface prior to plowing and,
2. Plow with a moldboard plow to a depth sufficient to mix the upper 8 inches of soil.

ALFALFA AND SAINFOIN

Sencor® herbicide is recommended for use in alfalfa and sainfoin in the following areas:

1. Alfalfa and sainfoin (Including mixed stands with grasses) (all areas except California).
2. Alfalfa and sainfoin (Including mixed stands with grasses) (California only).
3. Alfalfa - Tank Mix Combination with Gramoxone (Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming, and the following California counties: Del Norte, Lassen, Modoc, Nevada, Plumas, Shasta, Sierra, and Siskiyou).
4. Alfalfa - Post Dormant Application of Sencor® Impregnated on Dry Fertilizer Only (Connecticut, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New Mexico, New York, Ohio, Oklahoma, Pennsylvania, South Dakota, Tennessee, Texas and Wisconsin).

Sencor® 4 is recommended for use in aerial or ground spray equipment as a broadcast surface application to established crops of alfalfa and sainfoin for the control of certain grass and broadleaf weeds.

APPLICATION: Refer to "General Information" in the front of this label for detailed information on the application of Sencor®. For information on applying Sencor® 4 in fluid or on dry fertilizer refer to the "Application Of Sencor® 4 In Fluid Fertilizers" or "Commercial Impregnation And Application Of Sencor® 4 On Dry Bulk Fertilizer" under the "General Information" section of this label.

SPECIAL PRECAUTIONS:

- Use Sencor® 4 only on established alfalfa and sainfoin. Do not apply Sencor® 4 after growth begins in the spring or before growth ceases in the fall, except as specified on this label.
- Do not graze or harvest within 28 days after application.
- For best weed control, apply Sencor® 4 when weeds are less than 2 inches tall or before weed foliage is 2 inches in diameter.
- Reduced weed control may occur when extended dry conditions follow application of Sencor® 4.
- Crop injury may occur when:
 1. Crop is under stress conditions such as diseases, insect infestations, poorly drained soils, drought or winter injury at time of application;
 2. Crop is treated within 12 months after seeding;
 3. There is excessive irrigation or rainfall immediately after application. Do not apply more than 1/2 inch of water in the first irrigation after Sencor® 4 is applied.

ALFALFA and SAINFOIN
(All Areas Except California)
BROADCAST APPLICATION INSTRUCTIONS

| CROP | SENCOR® 4 Pt/Acre |
|--|-------------------|
| Alfalfa and Sainfoin (Except California) | 1/2 to 2 |

Select the proper dosage according to weeds known to be and present in field to be treated. On loamy sand soils in Oregon and Washington, do not apply more than 1 pt of Sencor® 4 per acre.

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES:

Rates of 1 to 1-1/2 pt of Sencor® 4 per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa. Higher rates will severely reduce forage grass stands.

Sencor® 4 should not be used on sand soils. In areas west of the Rocky Mountains, avoid using Sencor® 4 on soils with calcareous surface area, high levels of lime or sodium, or a pH greater than 8.2.

| ALFALFA and SAINFOIN Weeds Controlled (Except California) | | |
|---|--|--|
| 1/2 to 3/4 Pt SENCOR® 4/Acre | | |
| Chickweed, Common (<i>Stellaria media</i>) | | |
| 3/4 to 1 Pt SENCOR® 4/Acre | | |
| Cheat (<i>Bromus secalinus</i>) Deadnettle, Purple (<i>Lamium purpureum</i>) Downy brome (<i>Bromus tectorum</i>) | Japanese brome (<i>Bromus japonicus</i>) Pennycress (<i>Thlaspi arvense</i>) | Rescuegrass (<i>Bromus catharticus</i>) Shepherdspurse (<i>Capsella bursa pastoris</i>) |
| 1 to 2 Pt SENCOR® 4/Acre | | |
| Broadleaves Fleabane, Rough (<i>Erigeron strigosus</i>) Flixweed (<i>Descurainia sophia</i>) Henbit (<i>Lamium amplexicaule</i>) Kochia (<i>Kochia scoparia</i>) Lambsquarters, Common (<i>Chenopodium album</i>) | Marestail (Horseweed) (<i>Hippuris vulgaris</i>) Meadow Salsify (<i>Tragopogon pratensis</i>) Mustard, Blue (<i>Chorispora tenella</i>) Mustard, Jim Hill (tumble) (<i>Sisymbrium altissimum</i>) Mustard, Tansy (<i>Descurainia pinnata</i>) Pepperweed (<i>Lepidium virginicum</i>) | Pigweed, Redroot (<i>Amaranthus retroflexus</i>) Prickly Lettuce (<i>Lactuca serriola</i>) White Cockle (<i>Melandrium album</i>) Wild Buckwheat (<i>Polygonum convolvulus</i>) Yellow Rocket (<i>Barbarea vulgaris</i>) |
| Grasses Foxtail, Green (<i>Setaria viridis</i>) | Little Barley (<i>Hordeum pusillum</i>) Smooth Brome (<i>Bromus inermis</i>) | Wild Oats (<i>Avena fatua</i>) |
| 2 Pt SENCOR® 4/Acre | | |
| Broadleaves Chickweed, Mouseear (<i>Cerastium vulgatum</i>) | Dandelion (<i>Taraxacum officinale</i>) | Ragweed, Common (<i>Ambrosia artemisiifolia</i>) |
| Grasses Barnyardgrass (<i>Echinochloa crus-galli</i>) | Bluegrass (<i>Poa annua</i>) | Foxtail Barley (<i>Hordeum jubatum</i>) |

Weeds Partially Controlled: At the rate of 2 pt/acre Sencor® 4 may be used to reduce the competition from curly dock (*Rumex crispus*). At 1 to 2 pt/acre, Sencor® 4 may be used to reduce the competition of German Moss or knawel (*Scleanthus annus*).

ALFALFA AND SAINFOIN (California Only)
(Including Mixed Stands With Grasses)

Sencor® 4 is recommended for use in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin.

APPLICATION: Sencor® 4 is recommended for use in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply Sencor® 4 after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.

For information on applying Sencor® 4 in fluid fertilizer solutions to alfalfa, refer to the appropriate section of this label.

For information on commercial impregnation and application of Sencor® 4 on dry bulk fertilizer, refer to the appropriate section of this label.

| Weeds Controlled | | |
|--|---|--|
| 3/4 to 1 Pt Sencor® 4/Acre | | |
| Cheatgrass (downy brome) <i>(Bromus secalinus)</i> | | |
| 1 to 2 Pt Sencor® 4/Acre | | |
| Broadleaves Chickweed, Common (<i>Stellaria media</i>) Flixweed (<i>Descurainia sophia</i>) Henbit (<i>Lamium amplexicaule</i>) Kochia (<i>Kochia scoparia</i>) | Meadow Salsify (<i>Tragopogon pratensis</i>) Mustard, Blue (<i>Chlorispora tenella</i>) Mustard, Tansy (<i>Descurainia pinnata</i>) Pepperweed, Virginia (<i>Lepidium virginicum</i>) | Shepherdspurse (<i>Capsella bursa-pastoris</i>) White Cockle (<i>Melandrium album</i>) Wild Buckwheat (<i>Polygonum convolvulus</i>) Yellow Rocket (<i>Barbarea vulgaris</i>) |
| Grasses Smooth Brome (<i>Stellaria media</i>) | Wild Oats (<i>Avena fatua</i>) | |
| 2 Pt Sencor® 4/Acre | | |
| Broadleaves Dandelion (<i>Taraxacum officinale</i>) | | |
| Grasses Barnyardgrass (<i>Echinochloa crus-galli</i>) | Bluegrass (<i>Poa annua</i>) | Foxtail Barley (<i>Hordeum jubatum</i>) |

| BROADCAST APPLICATION INSTRUCTIONS | |
|---|--------------------------|
| CROP | Sencor® 4 Pt/Acre |
| Alfalfa and Sainfoin (California Only) | 3/4 to 2 |
| <p>Select the proper dosage according to weeds known to be present in the field to be treated. Apply specified dosage in 20 to 40 gallons of water per acre with ground spray equipment or 3 to 10 gallons of water per acre with aerial spray equipment fitted with nozzles suitable for broadcast applications of herbicides. Treat only dormant established crops of alfalfa and sainfoin. Injury may occur to alfalfa if Sencor® 4 is applied earlier than 12 months after seeding. Do not apply after Spring growth begins or before growth ceases in the Fall. Do not graze or harvest within 28 days after application.</p> <p>At the 2 pt/acre rate, Sencor® 4 may be used for suppression of curly dock.</p> | |

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES: Rates of 1 to 1-1/2 pt of Sencor® 4 per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa. Higher rates will severely reduce forage grass stands.

Do not apply with aerial spray equipment when wind speed is greater than 10 mph. Do not apply when weather conditions favor spray drift and/or when sensitive cool season crops, such as cole crops, onions, peas, or strawberries, are present in adjacent fields. Applications should not be made when weather conditions favor spray drift, especially in areas where wheat is growing on coarse textured soils in adjacent fields, or injury may occur.


ALFALFA

SENCOR® PLUS GRAMOXONE EXTRA TANK MIX

Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming and the following California counties: Del Norte, Lassen, Modoc, Nevada, Plumas, Shasta, Sierra, and Siskiyou.

APPLICATION: Sencor® 4 plus Gramoxone Extra herbicide tank mix application is recommended for use, during the dormant season, in aerial or ground spray equipment as a broadcast surface application to established alfalfa for the control of certain grass and broadleaf weeds. Do not apply Sencor®/Gramoxone Extra tank mix to growth that is more than 2 inches tall. Apply once per season. Do not apply following cuttings during growing season. Use a minimum of 10 gallons of water per acre with aerial spray equipment and a minimum of 20 gallons of water per acre with ground spray equipment. Add a non-ionic spreader at label rates to the spray solution.

Weeds Controlled: Sencor® 4 plus Gramoxone Extra (1-1/2 to 2-1/2 pints/acre) tank mix application will control established weeds. Gramoxone controls weeds by contact activity.



1/2 to 3/4 Pt of Sencor® 4/Acre

| | | | |
|------------------|--|--|--|
| Common Chickweed | | | |
|------------------|--|--|--|

3/4 to 1-1/2 Pt of Sencor® 4/Acre

| | | | |
|-----------|------------------|----------------|----------------|
| Bluegrass | Downy brome | Henbit | Rescuegrass |
| Cheat | Field pennycress | Japanese brome | Shepherdspurse |

Use Sencor® 4 at 1 to 1-1/2 Pt/Acre for control of the following weeds:

| | | | |
|----------------------|-----------------------|-----------------|----------------|
| Blue mustard | Kochia | Redroot pigweed | Tansy mustard |
| Common lambsquarters | Little barley | Rough fleabane | White cockle |
| Flixweed | Marestail (Horseweed) | Ryegrass | Wild oats |
| Green foxtail | Meadow salsify | Smooth brome | Wild buckwheat |
| Groundsel | Pepperweed | Sow thistle | Yellow rocket |
| Jim Hill mustard | Prickly lettuce | | |

| APPLICATION INSTRUCTIONS | |
|--|--|
| DOSAGE/ACRE | |
| Sencor® 4 1/2 to 1-1/2 Pt Plus Gramoxone Extra 1-1/2 to 2-1/2 Pt | Apply specified dosages of Sencor® and Gramoxone Extra in at least 10 gallons of water per acre with aerial equipment or at least 20 gallons of water per acre with ground equipment. Do not apply this tank mix to alfalfa growth if more than 2 inches tall. For best weed control, apply when broadleaf weeds and grasses are 1-6 inches tall and are actively growing. Care should be taken to avoid overlaps. Do not apply more than 1 pt of Sencor® 4 per acre on loamy sand soils. Reduced weed control may occur when extended dry conditions follow application of Sencor®. Crop injury may occur if alfalfa is under stress conditions such as diseases, insect infestations, drought or winter injury or if Sencor® is applied to alfalfa earlier than 12 months after seeding. |

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES: Rates of 1 to 1-1/2 pt of Sencor® 4 per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa.

Do not graze or harvest within 42 days after application.

In areas west of the Rockies, avoid the use of Sencor® on soils with calcareous surface, soils with high levels of lime or sodium, and with a pH greater than 8.2.

Do not apply when weather conditions favor spray drift. Aerial application should not be made when wind speed is greater than 10 mph. Do not use on sand soil.

Refer to the Gramoxone Extra label for additional directions, weed species controlled and precautions.

Post Dormant Application of Sencor® Impregnated on Dry Fertilizer Only

Sencor® 4 may be applied after dormancy has broken, but prior to three inches of new alfalfa shoot growth, only when impregnated on dry fertilizer in Connecticut, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New Mexico, New York, Ohio, Oklahoma, Pennsylvania, South Dakota, Tennessee, Texas and Wisconsin.

Apply at rates of 1-1/2 to 2 pt per acre as directed on this label for application during dormancy. Apply only when alfalfa foliage is dry or crop injury may occur. When using this application method, do not harvest or graze treated alfalfa for 60 days after application.

ASPARAGUS (Established)

Sencor® 4 is recommended for use in ground spray equipment or sprinkler irrigation (center pivot, lateral move, or solid set) systems as a single preemergence broadcast application or as a split application consisting of a preemergence broadcast application followed by a post harvest broadcast application.

Refer to the "General Information" section of this label for directions.

Weeds Controlled: Sencor® 4, applied to established asparagus according to directions, will effectively control:

| Broadleaves | | |
|--|--|--|
| Chickweed, Common (<i>Stellaria media</i>) | Pigweed, Redroot (<i>Amaranthus retroflexus</i>) | Smartweed, Pennsylvania (<i>Polygonum pensylvanicum</i>) |
| Jimsonweed (<i>Datura stramonium</i>) | Ragweed, Common (<i>Ambrosia artemisiifolia</i>) | Sorrel, Red (<i>Rumex acetosella</i>) |
| Lambsquarters (<i>Chenopodium album</i>) | | Velvetleaf (<i>Abutilon theophrasti</i>) |
| Grasses | | |
| Crabgrass (<i>Digitaria spp.</i>) | Foxtails (<i>Setaria spp.</i>) | Sandbur, Field (<i>Conchus pauciflorus</i>) |

| BROADCAST APPLICATION INSTRUCTIONS | |
|--|---|
| CROP | SENCOR® 4 Pt/Acre |
| Asparagus | 2 to 4 |
| | PREEMERGENCE APPLICATION ONLY Make a single surface application in early Spring before asparagus spears or ferns emerge. If the field is to be disked, apply Sencor® 4 after disking but before the crop emerges. Use the lower rate for control of the broadleaf weeds listed above. Use the higher rate in fields with a history of severe infestations of grasses and for maximum residual control. Do not apply within 14 days of harvest. |
| | 1 to 2 preemergence plus 2 to 3 post harvest |
| | SPLIT APPLICATION PREEMERGENCE AND POST HARVEST Preemergence Application: Apply before asparagus spears or ferns emerge. If the field is to be disked, apply after disking but prior to crop emergence. Do not apply within 14 days of harvest. Post Harvest Application: Apply after last harvest of the season but prior to emergence. The lower combination rates may be used for control of common ragweed, lambsquarters, redroot pigweed, and red sorrel. Use the higher combination rates for other weeds listed or in fields with severe grass infestations or for maximum post harvest control of emerged weeds. |
| IMPORTANT: The total amount of Sencor® 4 applied in one crop season may not exceed 4 pt per acre. | |

SPECIAL PRECAUTIONS (Asparagus):

- Do not use on newly seeded asparagus nor on young plants during the first growing season after setting crowns.
- DO NOT APPLY POST HARVEST APPLICATIONS UNTIL AFTER THE LAST HARVEST OF SPEARS.
- **Aerial application is prohibited.**

CARROTS

Special Conditions of Sale Provision for Use on Carrots: The following directions for use were developed under the direction of IR-4 (government minor crops use program). As such the testing was done independently from the testing program of Bayer CropScience. Buyer is advised that Bayer CropScience makes no assurances regarding satisfaction with the product and all risks of crop injury or product performance are assumed by the Buyer.

Apply Sencor® herbicide with ground equipment as specified below under "Application Instructions". For effective control of broadleaf weeds with postemergence applications, apply Sencor® before weeds are 1 inch in height or diameter. Thorough spray coverage is essential for adequate weed control.

Do not use air blast or other high pressure spray equipment to make postemergence applications of Sencor®. Refer to the appropriate section of this label for additional information regarding spray equipment, dilution rates, mixing, sprayer cleanup, restrictions, container disposal and cautions.

Refer to "Mixing" under the "General Information" section on the front of this label.

For specific application information see "General Information" and "Application" sections at the front of this label.

Weeds Controlled: Sencor® 4 applied to carrots according to directions will effectively control:

| | | |
|--|--|--|
| Carpetweed (<i>Mullugo verticillata</i>) | Mustard, Wild (<i>Sinapis arvensis</i>) | Prickly Lettuce (<i>Lactuca serriola</i>) |
| Galinsoga (<i>Galinsoga parviflora</i>) | Pigweed, Redroot (<i>Amaranthus retroflexus</i>) | Shepherdspurse (<i>Capsella bursa-pastoris</i>) |
| Horseweed (<i>Conyza canadensis</i>) | Pigweed, Smooth (<i>Amaranthus hybridus</i>) | Pineappleweed (<i>Matricaria matricarioides</i>) |
| Lambsquarters, Common (<i>Chenopodium album</i>) | | |

| APPLICATION INSTRUCTIONS | |
|---|--|
| CROP | SENCOR® 4 Pt/Acre |
| Carrots | 1/2 |
| | Apply specified dosage per acre as a broadcast spray over the tops of carrot plants. Application should be made after carrots have formed 5-6 true leaves but before weeds are 1 inch in height or diameter. If needed, a second application may be made after an interval of at least 3 weeks. Applications may be made up to 60 days of harvest. |
| IMPORTANT: The total amount of Sencor® 4 applied in one crop season must not exceed 1 pt per acre. | |

SPECIAL PRECAUTIONS:

- Do not apply to carrots grown for seed. Do not apply within 3 days after periods of cool, wet or cloudy weather or crop injury will occur. Do not apply Sencor® within 3 days of any other chemical unless specified on this label. Do not apply on very hot days or excessive crop injury will result. Do not apply until carrots have at least 5-6 true leaves. Earlier applications will result in excessive crop damage.
- Crop injury or delayed maturity may result from applications of Sencor® if carrots are growing under stress conditions such as periods of drought or cool, wet and cloudy weather preceding application.
- Following an application of Sencor® 4, chlorosis (yellowing) and burning of the leaf tissue may occur.
- For newly introduced varieties of carrots with unknown tolerance to Sencor®, treat only a small area to determine if Sencor® can be used without injury to the crop.

FIELD CORN

POSTEMERGENCE APPLICATION

Sencor® 4 is recommended for control of selected broadleaf weeds when applied as a tank mix combination with certain broadleaf herbicides presently registered and recommended for postemergence use in field corn. Herbicides which may be tank mixed with Sencor® 4 include:

| | | | |
|----------|---------------------|-----------------------------|--------------|
| 2,4-D | Buctril/Buctril Gel | Buctril + atrazine (Premix) | Resource |
| Atrazine | Laddok S-12 | Clarity | Scorpion III |
| Banvel | Marksman | Pursuit* | Tough |
| Basagran | | | |

* Use only on Pursuit resistant/tolerant corn hybrids (IMI- Corn).

APPLICATION: Sencor® 4 may be applied to field corn after crop emergence until just prior to tasseling. Broadcast applications may be made with ground or aerial equipment. For optimum weed control, apply treatments when weeds are small and actively growing, but before reaching the maximum heights listed in the Weeds Controlled table.

POSTEMERGENCE BROADCAST APPLICATION

Ground Application: Adjust nozzle height above crop and weed canopy to ensure uniform spray coverage. Gallonage should be increased with increasing weed size and population density.

For tank mixes of Sencor® 4 plus atrazine, Basagran, Laddok S-12, Buctril, Buctril + atrazine, Pursuit, Resource, Tough or 2,4-D amine formulations, use flat fan nozzles spaced a maximum of 20 inches apart. Best results are achieved using a minimum spray volume of 10 gallons per acre and spray pressure from 20 to 40 psi.

For Sencor® 4 tank mixes with Banvel, Clarity, Marksman, or 2,4-D low volatile ester formulations, use drift-reducing nozzles which are specifically designed to produce coarse sprays and reduce the amount of driftable fines. Additional measures which will help avoid potential drift to sensitive crops and plants include using a minimum spray volume of 20 gallons per acre and keeping spray pressures at or below 20 psi unless otherwise specified by the nozzle manufacturer.

For further precautions and additional instructions, consult the tank mix partner's label.

Aerial Application: Apply in a minimum spray volume of 3 gallons per acre. For optimum spray coverage and distribution, use a minimum of 5 gallons per acre and a maximum pressure of 40 psi. Use a boom and nozzle configuration which will provide a uniform deposition pattern and coverage with low drift potential. Avoid overlaps to prevent potential crop injury. Do not apply near sensitive crops or sensitive plants growing near the treated area. Do not apply when wind speed is greater than 10 mph or when winds are moving toward sensitive crops or plants. To avoid drift hazards, applicator must follow the most restrictive labeling of the products used in a tank mix. Refer to the appropriate tank mix partner's label for further precautions and instructions.

POST DIRECTED APPLICATION

Sencor® 4 in tank mix combinations with Banvel, 2,4-D, Buctril, or Scorpion III may be applied post directed to field corn. Use drop nozzles and appropriate spacing to direct spray below the corn whorl and upper leaves. The top of the target weed canopy must be sufficiently below the whorl and upper leaves of the crop to permit this application and provide adequate spray coverage. The height differential required between the crop and weed canopy will depend on the specific equipment used. Apply before tassel emergence. For further precautions and additional instructions, refer to the appropriate tank mix partner's label.

ADJUVANTS

The adjuvant types listed below may be utilized with certain Sencor® 4 tank mix combinations. Consult the tank mix instructions section for the appropriate adjuvant and rate. Use of non-recommended adjuvants or rates may result in severe leaf burn, crop stunting, and/or stand reduction. Use only adjuvants which are exempt from tolerance requirements under 40 CFR 180.1001.

UAN (urea ammonium nitrate) is commonly referred to as 28, 30, or 32% N.

Ammonium sulfate (spray grade) may be used as an alternative to UAN with certain tank mix combinations.

Non-ionic surfactants should contain at least 80% active ingredient.

DO NOT USE crop oil concentrate (COC) or any adjuvant containing vegetable or petroleum oils with any Sencor® 4 tank mixtures as severe leaf burn, crop stunting, and/or stand reduction may occur.

RAINFASTNESS

Sencor® 4 will not reduce rainfastness of the recommended tank mix partners. Refer to the individual product labels for rainfastness recommendations.

SPRAYER CLEANUP

Refer to each tank mix partner's label and the Sprayer Cleanup section of the Sencor® 4 label for specific instructions on cleaning spray equipment. Special attention should be given to the required cleanup procedures for 2,4-D, Banvel, Clarity, and Marksman.

SPECIAL PRECAUTIONS

- Do not use on corn grown for seed, sweet corn, popcorn, or white corn.
- Do not apply more than 0.25 pounds a.i. metribuzin (8 fl oz Sencor® 4) per acre per use season.
- Do not apply when field corn is under stress (see Stress statement below).
- Do not use aerial applications if sensitive crops or plants are growing in the vicinity of the area to be treated.
- Do not allow spray drift onto sensitive crops or plants.
- Do not use on sand, loamy sand or sandy loam soils that have less than 0.5% organic matter.
- Do not use on sand or loamy sand soils in Washington, Oregon or Idaho or crop injury may occur.
- Observe all precautions and limitations on labeling of all products used in the tank mixtures.

Stress is any condition or combination of conditions which impairs normal crop growth. Weather, disease, insect damage, fertility or other factors may cause stress. Applications made before or after the corn is under stress from these factors or from periods of prolonged cool, wet and cloudy weather or widely fluctuating day and nighttime temperatures, may result in temporary leaf burn, yellowing and/or stunting of the crop. Recovery from damage is generally rapid with no lasting effects on new growth. Under extreme stress, stand reductions may occur.

Feeding Restrictions: Field corn treated with Sencor® 4 may be grazed or harvested for silage or grain 60 days after treatment. Follow the most restrictive preharvest interval on the labels of the products used in the tank mixtures.

TANK MIX COMBINATIONS

The Sencor® 4 tank mixtures listed below can be utilized for control of certain annual broadleaf weeds.

| SENCOR® 4 POST DIRECTED INSTRUCTIONS | | |
|---|---|---|
| PRODUCT | RATE | DIRECTIONS AND REMARKS* |
| Sencor® 4 + 2,4-D Amine or 2,4-D LVE | 3 to 4.5 fl oz/A + 3/4 to 1-1/2 pt/A ¹ or 1/2 to 3/4 pt/A ¹ | For corn greater than 8-inches tall, apply as a directed spray with drop nozzles before tassel emergence. Apply only to varieties known to be tolerant to 2,4-D. DO NOT USE ADJUVANTS. 2,4-D may cause injury to nearby sensitive crops. 2,4-D applications may result in brittle corn stalks, and winds or cultivation may cause stalk breakage. To reduce damage, delay cultivation 8 to 10 days after application. |
| Sencor® 4 + Banvel | 3 to 4.5 fl oz/A + 1/2 pt/A | For corn 8 to 36 inches tall, apply as a directed spray with drop nozzles . Application may be made up to 15 days prior to corn tasseling. If growing conditions are dry and plants are stressed, addition of a non-ionic surfactant (1 qt/100 gal of spray solution) may improve weed control. For corn grown on coarse textured soils, apply Banvel at 0.5 pt/acre, regardless of application method. Application may cause injury to nearby sensitive crops or plants. Application may result in temporary leaning of corn plants. Delay cultivation until plants return to normal growth patterns to avoid stalk breakage. |
| Sencor® 4 + Buctril or Buctril Gel | 3 to 4.5 fl oz/A + 1 to 1-1/2 pt/A or 1/2 to 3/4 pt/A | Apply as a directed spray with drop nozzles before tassel emergence. DO NOT USE ADJUVANTS. Occasional temporary corn leaf burn may occur and is similar to that observed from liquid fertilizers. Recovery is generally rapid with no lasting effect. To reduce potential for crop damage, application should be made to dry corn foliage when weather conditions are not extreme. |
| Sencor® 4 + Scorpion III | 3 to 4.5 fl oz/A + 4 oz/A | For corn 8 to 24 inches tall, apply as a directed spray with drop nozzles. Include nonionic surfactant (1 qt/100 gal) plus UAN (2.5 gal/100 gal) for optimum weed control. |
| * Consult the appropriate tank mix partner's label for additional instructions or restrictions. The most restrictive labeling applies to tank mixes with Sencor® 4. | | |
| ¹ Application rate is based on, but not restricted to, 4 pounds active ingredient per gallon of 2,4-D. | | |

SENCOR® 4 POSTEMERGENCE BROADCAST INSTRUCTIONS

| PRODUCT | RATE | DIRECTIONS AND REMARKS* |
|---|--|---|
| Sencor® 4 + 2,4-D Amine or 2,4-D LVE | 3 fl oz/A + 1/2 to 1 pt/A ¹ or 1/3 to 1/2 pt/A ¹ | Apply as a broadcast spray during the interval from corn emergence until corn is 8 inches tall. Apply only to varieties known to be tolerant to 2,4-D. DO NOT USE ADJUVANTS. 2,4-D may cause injury to nearby sensitive crops. 2,4-D applications may result in brittle corn stalks, and winds or cultivation may cause stalk breakage. To reduce damage, delay cultivation 8 to 10 days after application. |
| Sencor® 4 + Atrazine | 3 fl oz/A + 1/2 to 1-1/2 lb ai/A | Apply as a broadcast spray during the interval from corn emergence until corn is 12 inches tall. A non-ionic surfactant (1 qt/100 gal of spray solution) may be added to improve weed control. Atrazine is a restricted use herbicide. Follow all state and federal label instructions and restrictions pertaining to atrazine applications. |
| Sencor® 4 + Banvel or Clarity | 3 fl oz/A + 1/2 to 1 pt/A or 1/2 to 1 pt/A | Apply as a broadcast spray during the interval from corn emergence through the five leaf stage or when corn is 8 inches tall, whichever occurs first. For Banvel applications to corn greater than 8 inches in height, consult the Banvel label for use rates and restrictions. If growing conditions are dry and plants are stressed, addition of a non-ionic surfactant (1 qt/100 gal of spray solution) may improve weed control. For corn grown on coarse textured soils, apply Banvel or Clarity at 0.5 pt/A, regardless of application method. Application may cause injury to nearby sensitive crops or plants. Application may result in temporary leaning of corn plants. Delay cultivation until plants return to normal growth patterns to avoid stalk breakage. |
| Sencor® 4 + Basagran | 3 fl oz/A + 1 pt/A | Apply as a broadcast spray after corn emergence but before corn exceeds 30 inches in height and the crop canopy closes the row. Adjuvants such as UAN (0.5 to 1 gal/A), ammonium sulfate (17 lbs/100 gal of spray solution), or non-ionic surfactant (1 qt/100 gal of spray solution) may improve weed control. |
| Sencor® 4 + Buctril or Buctril Gel | 2.4 to 3 fl oz/A + 1 pt/A or 1/2 pt/A | Apply as a broadcast spray when corn is in the fourth true leaf stage or later but before the crop canopy closes the row. DO NOT USE ADJUVANTS. Occasional temporary corn leaf burn may occur and is similar to that observed from liquid fertilizers. Recovery is generally rapid with no lasting effect. To reduce potential for crop damage, application should be made to dry corn foliage when weather conditions are not extreme. |
| Sencor® 4 + Buctril + atrazine (Premix) | 2.4 to 3 fl oz/A + 1-1/2 to 2 pt/A | Apply as a broadcast spray during the interval from corn emergence until corn is 12 inches tall. DO NOT USE ADJUVANTS. Occasional temporary corn leaf burn may occur and is similar to that observed from liquid fertilizers. Recovery is generally rapid with no lasting effect. To reduce potential for crop damage, application should be made to dry corn foliage when weather conditions are not extreme. |
| Sencor® 4 + Laddok S-12 | 3 fl oz/A + 1.33 to 1.66 pt/A | Apply as a broadcast spray after corn emergence until the corn is 12 inches tall. Adjuvants such as UAN (0.5 to 1 gal/A) may improve weed control. Laddok S-12 contains atrazine, and is a restricted use product. Follow all state and federal label instructions and restrictions pertaining to atrazine. |
| Sencor® 4 + Marksman | 3 fl oz/A + 1-1/2 to 2 pt/A | Apply as a broadcast spray during the interval from corn emergence through the five-leaf stage or when corn is 8 inches tall, whichever occurs first. DO NOT USE ADJUVANTS. Application may cause injury to nearby sensitive crops or plants. Application may result in temporary leaning of corn plants. Delay cultivation until plants return to normal growth patterns to avoid stalk breakage. Marksman contains atrazine, and is a restricted use product. Follow all state and federal label instructions and restrictions pertaining to atrazine. |
| Sencor® 4 + Pursuit | 3 fl oz/A + 2 to 4 oz/A | Use only on designated IMI-Corn hybrids (hybrids which are resistant/ tolerant to Pursuit). Apply the 4.0 ounce rate of Pursuit if grasses are present or broadleaf weeds are near the maximum heights shown. Apply in combination with a non-ionic surfactant (1 qt/100 gal of spray solution) and UAN (1 to 2 qt/A). |
| Sencor® 4 + Resource | 3 fl oz/A + 4 to 6 fl oz/A | Apply as a broadcast spray to field corn from 2-leaf through 10-leaf (visible leaf collars) stage. Adjuvants such as nonionic surfactant (0.25% v/v), UAN (2% v/v) or ammonium sulfate (2.5 lbs/A) may improve weed control. |
| Sencor® 4 + Tough | 3 fl oz/A + 1 to 2 pt/A | Apply as a broadcast spray after corn emergence but before corn height exceeds 30 inches and the crop canopy closes the row. A non-ionic surfactant (1 qt/100 gal of spray solution) may be added to improve weed control. Use the higher rates of Tough as weeds approach the maximum height listed or are found in high density. Tough may improve control on triazine/ALS resistant weeds. |

* Consult the appropriate tank mix partner's label for additional instructions or restrictions. The most restrictive labeling applies to tank mixes with Sencor® 4.

¹ Application rate is based on, but not restricted to, 4 pounds active ingredient per gallon of 2,4-D.

| WEEDS CONTROLLED - POST DIRECTED APPLICATION | | | | |
|---|---------------------------------|--------|---------|--------------|
| These tank mixtures with Sencor® 4 will control the following annual weeds up to the maximum weed heights listed: | | | | |
| COMMON WEED NAME | Sencor® 4 + | | | |
| | 2,4-D | Banvel | Buctril | Scorpion III |
| | MAXIMUM WEED HEIGHT IN INCHES * | | | |
| Amaranth, Palmer | 12 | 12 | 6 | 8 |
| Cocklebur, common | 12 | 12 | 12 | 15 |
| Jimsonweed | 12 | 10 | 10 | 8 |
| Ladysthumb | 6 | 8 | 6 | 6 |
| Lambsquarters, common | 12 | 12 | 10 | 12 |
| Morningglory, entire leaf | 18 | 18 | 6 | 12 |
| Morningglory, ivyleaf | 18 | 18 | 6 | 12 |
| Morningglory, pitted | 18 | 18 | 6 | 12 |
| Morningglory, tall | 18 | 18 | 6 | 12 |
| Nightshade, black | 10 | 8 | 8 | 6 |
| Nightshade, eastern black | 10 | 8 | 8 | 6 |
| Pigweed, redroot | 12 | 12 | 6 | 8 |
| Pigweed, smooth | 12 | 12 | 6 | 8 |
| Ragweed, common | 8 | 8 | 8 | 10 |
| Ragweed, giant | 12 | 12 | 8 | 15 |
| Smartweed, Pennsylvania | 6 | 8 | 6 | 6 |
| Sunflower, common | 12 | 12 | 12 | 12 |
| Velvetleaf | 10 | 8 | 8 | 8 |
| Waterhemp, tall | 12 | 12 | 6 | 8 |

* When weeds are approaching the maximum height listed or found in high densities, use the higher rate of Sencor® 4 and the selected tank mix partners.

PERENNIAL WEED SUPPRESSION

The following Sencor® 4 tank mixtures will provide top growth burndown and in season suppression of the following perennial weeds; however, regrowth may occur. For the best performance on these weeds, use the maximum rates of Sencor® 4, Banvel, Buctril, Buctril + atrazine, Clarity, Marksman, 2,4-D LVE or Pursuit specified for these tank mixtures.

Sencor® 4 + Banvel or Clarity

Bindweed, field; Dandelion, common; Dock, curly; Smartweed, swamp; Thistle, Canada.

Sencor® 4 + Buctril or Buctril + atrazine

Thistle, Canada.

Sencor® 4 + 2,4-D LVE


Bindweed, field; Dandelion, common; Dock, curly; Smartweed, swamp; Thistle, Canada.

Sencor® 4 + Marksman

Bindweed, field; Dandelion, common; Dock, curly; Smartweed, swamp; Thistle, Canada.

Sencor® 4 + Pursuit

Thistle, Canada.



PREPLANT AND PREEMERGENCE APPLICATIONS

(Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota and Wisconsin)

Sencor® 4 is recommended for additional residual control of certain broadleaf weed species in corn when applied as a tank mix combination with both grass and broadleaf herbicides registered and recommended for use in field corn. Sencor® 4 can be tank mixed with specified rates of the following herbicides:

| | | | |
|--------------------|--------------|---------------|-----------------|
| Alachlor | Bullet | Harness Xtra | Pursuit* |
| Atrazine | Clarity | Lariat | Ramrod |
| Banvel | Cycle | Linex | Ramrod/Atrazine |
| Bicep | Dual | Linuron | Simazine |
| Bicep II | Dual II | Lorox | Surpass |
| Bicep Lite | Extrazine II | Marksman | Surpass 100 |
| Bladex | Frontier | Pentagon | Topnotch |
| Broadstrike + Dual | Guardsman | Prowl | |
| Broadstrike Plus | Harness | Pursuit Plus* | |

* Use only on Pursuit resistant/tolerant corn hybrids (IMI corn).

Application: Sencor® 4 may be applied to field corn preplant without incorporation up to 30 days prior to planting or preemergence. Applications may be made by either ground or aerial equipment. For tank mixes, follow the most restrictive application methods of all products used.

SPECIAL PRECAUTIONS:


- Do not apply more than 8 fluid ounces Sencor® 4 (0.25 pound active ingredient) per acre per growing season.
- Do not apply on soils having pH 7.0 or greater.
- Corn seed should be planted a minimum of 1-1/2 inches deep.
- Sencor® 4 may only be used in hybrid seed corn production fields if both inbred parents are known to be tolerant to Sencor® 4.
- Do not use on muck soils as reduced weed control may result.
- Observe all precautions and limitations on labeling of all products used in tank mixes.

Feeding restrictions: Corn treated with Sencor® 4 may be harvested for silage or grain 60 days after treatment. For tank mixes, follow the most restrictive preharvest interval of all products used.

Weeds Controlled*: Sencor® 4 will aid in the residual preemergence control of the following weed species when tank mixed with other registered grass and/or broadleaf corn herbicides:

| | | | |
|-----------------------|-----------------|-------------------------|-----------------|
| Horseweed/marestail | Pigweed spp. | Smartweed, Pennsylvania | Velvetleaf |
| Ladysthumb | Ragweed, common | Sunflower | Waterhemp, Tall |
| Lambsquarters, common | | | |

* For control of emerged weeds refer to the "Burndown Weed Control" section of the Sencor® label.



| SENCOR® 4 FIELD CORN RATE INSTRUCTIONS | | | |
|---|--------------------------|--|-----------------------|
| STATES | | APPLICATION TIMING | SENCOR® 4 fl ozs/A |
| Iowa Kansas Missouri | Nebraska South Dakota | Preplant (0 to 30 days) Preemergence | 3 to 8 |
| Illinois Indiana Kentucky Michigan | Minnesota Ohio | Preplant (10 to 30 days) | 3 to 8 |
| | Wisconsin | Preplant (0 to 9 days) Preemergence | 3 to 6 |

REMARKS: Apply as a broadcast spray prior to corn emergence from the soil.
 Do not apply Sencor® 4 on coarse textured soils with less than 1.5% organic matter.
 Do not apply more than 6 fl. oz. Sencor® 4 per acre on soils with less than 2.0% organic matter.
 For heavy weed infestations and/or early preplant applications, use the higher rates of Sencor® 4.
 Consult the label of herbicide tank mix partners to determine proper use rates for the other product(s).

BURNDOWN WEED CONTROL - FIELD CORN See page 26

GARBANZO BEANS (Chickpeas)

California, Idaho, Oregon, and Washington

Special Conditions of Sale for Use on Garbanzo Beans (Chickpeas): The following directions for use were developed under the direction of IR-4 (government minor crops use program). As such the testing was done independently from the testing program of Bayer CropScience. Buyer is advised that Bayer CropScience makes no assurances regarding satisfaction with the product and that all risks of crop injury or product performance are assumed by the Buyer.

Sencor® herbicide is recommended as a preemergence application for the suppression of certain broadleaf weeds in garbanzo beans.

Weeds Suppressed*:

| | | |
|----------------------|----------------------|--------------|
| Common Chickweed | Henbit | Wild Mustard |
| Dog Fennel (Mayweed) | Common Lambsquarters | Pigweed |
| Field Pennycress | Shepherdspurse | |

* Suppression is a reduction in weed size and growth compared to a non-treated area in the same field. Sencor® used alone will not control triazine-resistant weed species.

| APPLICATION INSTRUCTIONS | |
|--------------------------|---|
| CROP | SENCOR® 4 Pt/Acre |
| Garbanzo beans | 1/2 to 3/4 |
| | <p>Apply specified dosage in a single preemergence application using 10 to 40 gallons of water per acre with ground spray equipment. Apply before or after planting but before crop emergence. Thorough incorporation, either by rainfall or by mechanical means, is essential for weed suppression. Under dry conditions, incorporate Sencor® into the top 1 to 2 inches of soil with spike harrows, or similar shallow incorporation equipment, then cross harrow to insure uniform soil incorporation. Where soil surface is moist at the time of application and rain follows before weed emergence, a broadcast application should provide adequate weed suppression.</p> <p>Use on coarse-textured soils, sandy soils or any soil with less than 1.5% organic matter will likely cause crop injury.</p> <p>Use the higher rate on fine textured soils (high in clay or organic matter) and in fields with a history of high weed populations.</p> |

SPECIAL PRECAUTIONS:

- Crop injury may result if crop is under stress conditions caused by cold weather, poor soil fertility, disease or insect damage.
- Crop injury may result if application is followed by heavy rain. Avoid application of more than 1/2 inch of irrigation within one month after application of Sencor®, or crop injury may occur.
- Do not use on clay knobs or poorly covered subsoils.
- Do not apply preemergence on shallow seedings less than 2 inches deep.
- Do not graze or feed treated vines to livestock within 40 days after application.
- Maintain continuous spray tank agitation to keep material in suspension. Avoid overlapping of spray swaths and shut off spray booms while turning, slowing or stopping, or crop injury will occur.

NOTE: This treatment may cause some chlorosis or minor necrosis. Because garbanzo bean varieties may vary in their susceptibility to Sencor®, determine crop tolerance prior to adoption as a field scale practice to prevent possible injury.

LENTILS AND PEAS
(Idaho, Oregon, and Washington)

Sencor® 4 herbicide is recommended as a preemergence and postemergence application for the suppression of certain broadleaf weeds in lentils and peas.

Weeds Suppressed*

| | | |
|--|------------------------|--------------------|
| Common Chickweed** | Henbit** | Prostrate Knotweed |
| Corn Spurry | Lambsquarters | Redroot Pigweed |
| Dog Fennel | Pennsylvania Smartweed | Shepherdspurse** |
| Field Pennycress | Pineapple Weed | Wild Mustard |
| * Suppression is a reduction in weed size and growth compared to a non-treated area in the same field. | | |
| ** Preemergence application only. | | |

PREEMERGENCE APPLICATION: Make a single preemergence application of Sencor® 4 at 3/8 to 3/4 pint per acre per crop year. Apply in 10 or more gallons of water per acre with ground spray equipment or 5 or more gallons of water per acre with aerial spray equipment. Apply Sencor® 4 before or after planting. Thorough incorporation, either by rainfall or by mechanical means, is essential for weed suppression. Under dry conditions, incorporate Sencor® 4 into the top 1 to 2 inches of soil with spike harrows, or similar shallow incorporation equipment, then cross harrow to ensure uniform soil incorporation. Where soil surface is moist at the time of application and rain follows before weed emergence, a broadcast application should provide adequate weed suppression.

Use the higher rate on fine-textured soils (high in clay or organic matter) and in fields with a history of high weed populations.

Sencor® 4 may be applied pre- or post plant incorporated as a tank mix combination with FARGO 4EC. Follow the Directions for Use statements on both product labels.

POSTEMERGENCE APPLICATION: One postemergence application may be made per season. Use 1/4 to 1/2 pint of Sencor® 4 per acre on **lentils** and **spring peas**. On **winter peas**, use 3/8 to 1/2 pint of Sencor® 4 per acre. For suppression of dog fennel, use 1/2 pint Sencor® 4 per acre. Apply specified dosage in 20 or more gallons of water per acre with ground spray equipment or 5 or more gallons of water per acre with aerial spray equipment. Do not exceed 40 psi with ground spray equipment. Apply as a broadcast spray when weeds are small (less than 2 inches in height or diameter) and before crop is 6 inches tall.

Temporary chlorosis of the crop may occur. There is an added risk of crop injury if a postemergence application is made following a previous preemergence or post plant incorporated Sencor® application.

Do not apply over very moist soils or wet crop foliage. Do not apply postemergence within 3 days after periods of cool, wet, or cloudy weather or crop injury may occur.

Do not apply within 24 hours of treatment with other pesticides.

SPECIAL PRECAUTIONS (all applications): Do not apply more than 1 pint of Sencor® 4 per acre per year. Crop injury may result if crop is under stress conditions caused by cold weather, low fertility, disease or insect damage.

Crop injury may also result if application is followed by heavy rain.

Do not use on coarse-textured soils, sandy soils or soils with less than 1.5% organic matter.

Do not apply to "Estin" lentils.


Do not use on clay knobs or poorly covered subsoils.

Do not apply on shallow seedlings less than 2 inches deep (preemergence only).

Do not apply within 50 days of harvest of peas, or within 75 days of harvest of lentils. Do not graze or feed treated vines to livestock within 40 days after application.

Maintain continuous spray tank agitation to keep material in suspension. Avoid overlapping and shut off spray booms while turning, slowing or stopping, or crop injury will occur.

NOTE: This treatment may cause some chlorosis or minor necrosis. Because lentil and pea varieties may vary in their susceptibility to Sencor® 4, determining crop tolerance prior to adoption as a field scale practice is suggested to prevent possible injury.



For additional precautions, restrictions, limitations, and sprayer clean-up information refer to the appropriate sections of this label.

SUGARCANE

SPECIAL PRECAUTIONS

- For aerial and chemigation application methods on sugarcane the maximum application rate is 4 pts Sencor® 4/acre.
- To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product by aircraft at a minimum upwind distance of 400 ft from sensitive plants.
- Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4. Refer to the *Crop Rotation Restrictions* section of this label for more information.

SUGARCANE (Hawaii Only)


Sencor®, a selective herbicide, is effective as a preemergence and an early postemergence broadcast application for control of certain grass and broadleaf weeds. When applied as a spot treatment, it also provides excellent control of perennial grasses and broadleaves.

Ground Application: Sencor® 4 should be mixed by filling the spray tank half full of clean water. Then add the specified amount of Sencor® 4 to suit the total tank capacity and the rate of application per acre (preferably 25 to 35 gallons per acre). Complete filling the tank and maintain sufficient agitation during mixing and spraying to insure a uniform spray mixture.

Aerial Application: Sencor® 4 is recommended for use in aerial spray equipment as a preemergence or postemergence application to irrigated sugarcane. Aerial spray equipment should be calibrated to apply the proper amount of Sencor® 4 in 7 to 10 gallons of spray mixture per acre.

Sencor® 4 applied preemergence or postemergence to the sugarcane as a broadcast spray or spot treatment will effectively control the following when weeds are less than 3 inches in height.

| Weeds Controlled in Irrigated and Non-irrigated Sugarcane | | | |
|---|--|--|--|
| Broadleaves Amaranth, Spiny (<i>Amaranthus spinosus</i>) | Euphorbia, Wild (<i>Euphorbia</i> spp.) Fireweed (<i>Erechtites hieracifolius</i>) | Floras paintbrush (<i>Emilia sonchifolia</i>) Spurge, Garden (<i>Euphorbia hirta</i>) | Spurge, Graceful (<i>Euphorbia glomerifera</i>) |
| Grasses Crabgrass (<i>Digitaria</i> spp.) | Guineagrass (<i>Panicum maximum</i>) Plushgrass (<i>Chloris radiata</i>) | Ricegrass (<i>Oryzopsis hymenoides</i>) | Wiregrass (<i>Eleusine indica</i>) |
| Weeds Controlled in Irrigated Sugarcane Only | | | |
| Broadleaves Amaranth, Spleen (<i>Amaranthus dubius</i>) Haole Koa (<i>Leucaena leucocephala</i>) | Hialoa (<i>Waltheria americana</i>) Hilahila (<i>Mimosa pudica</i>) | Purslane, Common (<i>Portulaca oleracea</i>) | Rattlepod (<i>Crotalaria spectabilis</i>) |
| Grasses Alexandergrass (<i>Brachiaria plantaginea</i>) | Bristly foxtail (<i>Setaria verticillata</i>) | | |
| Weeds Controlled in Non-irrigated Sugarcane Only | | | |
| Broadleaves Ageratum (<i>Ageratum conyzoides</i>) | Richardia (<i>Richardia brasiliensis</i>) | Tarweed (<i>Cuphea carthagenesis</i>) | |



| SUGARCANE (HAWAII ONLY) | |
|--|--|
| BROADCAST APPLICATION INSTRUCTIONS | |
| SENCOR® 4 Pt/Acre | REMARKS |
| 4 to 8 (1/2 to 1 gal) (non-irrigated) | PREEMERGENCE (Irrigated and non-irrigated sugarcane): Apply specified dosage per acre as a broadcast spray to the soil surface. Applications should be made within two weeks after planting prior to cane emergence or shortly after emergence (spike stage). OR EARLY POSTEMERGENCE (Irrigated and non-irrigated sugarcane): Apply specified dosage per acre as a broadcast spray over the cane. Application may be delayed as long as 4 to 6 weeks after planting provided weeds are less than 3 inches in height. |
| 8 to 12 (1 to 1-1/2 gal) (irrigated) | |
| 4 to 8 (1/2 to 1 gal) | OR POSTEMERGENCE: Apply specified dosage per acre as a broadcast spray to control weeds prior to "close in" time when cane shades out the weed growth. |
| 5 to 10 (1-1/8 to 1-1/4 gal) | SPOT TREATMENT: Apply specified dosage in 30 to 50 gallons of finished spray per acre. Spot treatments may be used to control weeds in missed areas, corners of fields, or areas of hard to control weeds. |
| NOTE: Do not apply more than 16 pt (2 gal) of Sencor® 4 (8 lb active ingredient)/ acre per crop cycle regardless of the method of application. The last application may be made up to 17 months of harvest. | |

SPECIAL PRECAUTIONS:

- Do not use treated foliage for feed or forage.

SUGARCANE (Louisiana and Texas Only)

| Preemergence and postemergence applications of Sencor® 4 with aerial or ground spray equipment are recommended for control of the following weeds in sugarcane in Louisiana and Texas: | | | |
|--|--|---|---|
| Broadleaves Amaranth, Spiny (Amaranthus spinosus) Bindweed, Field (Convolvulus arvensis) | Chickweed (Cerastium vulgatum) Henbit (Lamium amplexicaule) Lambsquarters (Chenopodium album) | London rocket (Sisymbrium irio) Marestail (Conyza canadensis) Mustard, Wild (Brassica kaber) | Pigweeds (Amaranthus spp.) Purslane (Portulaca oleracea) Sowthistle (Sonchus spp.) |
| Grasses Broadleaf Signalgrass (Brachiaria platyphylla) | Crabgrass (Digitaria spp.) Foxtails (Setaria spp.) | Johnsongrass, Seedling (Sorghum halepense) | Oats, Winter (Avena spp.) |

| SUGARCANE (LOUISIANA AND TEXAS ONLY) | |
|--------------------------------------|--|
| BROADCAST APPLICATION INSTRUCTIONS | |
| Sencor® 4 Pt/Acre | REMARKS |
| 3 to 6 | BROADCAST: Apply specified dosage per acre using 20 to 30 gallons of water with ground equipment or 5 gallons of water with aircraft spray equipment. Apply as a broadcast spray during the Fall after planting or to the stubble after harvest. Make a second application early in the Spring. |
| 1-1/2 to 3 | BAND: Apply specified dosage in 10 to 20 gallons of water per acre in a 30 to 36 inch band, over the row during the Fall after planting or to the stubble after harvest. Make a second application early in the Spring. |

SPECIAL PRECAUTIONS (Louisiana and Texas only):

- Do not use treated foliage for feed or forage.
- Use the higher rate on heavy clay soil and soil with a high percentage of organic matter. If necessary, a third application may be made in late Spring at layby. Do not apply within 60 days of harvest.
- For aerial and chemigation application methods on sugarcane the maximum application rate is 4 pts Sencor 4/acre.

- To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

SUGARCANE (Florida Only)

| | | | |
|---|---|--|---|
| Postemergence over-the-top or directed spray applications of Sencor® 4 are recommended for the control of the following weeds in sugarcane in Florida. | | | |
| Broadleaves Amaranth, Spiny (seedling) (Amaranthus spinosus) | Butterweed (Cressleaf groundsel) (Senecio glabellus) | Purslane (Portulaca oleracea) | |
| Grasses *Crabgrass, large (Digitaria sanguinalis) | Foxtail, bristlegrass (Setaria magna) Goosegrass (Eleusine indica) | Panicum, broadleaf (Panicum adspersum) | Signalgrass, Broadleaf (Brachiaria platyphylla) |
| *Best control is achieved when applications are made when this weed is less than 4" in diameter. | | | |

| SUGARCANE (FLORIDA ONLY) | |
|---------------------------------|--|
| APPLICATION INSTRUCTIONS | |
| Sencor® 4 Pt/Acre | REMARKS |
| 2 to 4 | GROUND APPLICATION: Sencor® 4 may be used in one or two applications with a minimum of 14 days between each application. Apply when weeds are less than 6 inches tall in 10 to 40 gallons of spray mixture per acre. POSTEMERGENCE BROADCAST OR BAND: Apply over the top of stubble or plant cane while sugarcane is less than 14 inches tall. POSTEMERGENCE DIRECTED SPRAY: Apply to sugarcane that is a minimum of 14 inches tall and before row closing. |
| 2 to 3 | AERIAL APPLICATION: Apply when weeds are less than 4 inches tall in 5 to 10 gallons of spray mixture per acre. Apply to stubble or plant cane while the sugarcane is less than 14 inches tall. |

SENCOR® PLUS ATRAZINE TANK MIX: Sencor® 4 may be used with atrazine as a preemergence or postemergence (before row closing) application to sugarcane. Rates for Sencor® 4 are 1-1/2 to 4 pt/acre and atrazine 80% W/P (4L) are 2-1/2 to 5 lb/acre (2 to 4 qt/acre). For additional information on precautions, instructions, limitations, application, and weeds controlled, refer to this label and the atrazine label.

SPECIAL PRECAUTIONS (Florida only): Do not use more than 4 pt per acre in a single growing season. Do not use on sand soils. Spray contact with sugarcane foliage may result in minor leaf margin chlorosis and/or necrosis. Do not apply within 60 days of harvest. Do not use treated crop for feed or forage. Avoid spray overlaps or variations in application speed that may result in insufficient or excessive rates of application.

To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product by aircraft at a minimum upwind distance of 400 ft from sensitive plants.


TOMATOES

Apply Sencor® herbicide with ground equipment to seeded and transplanted tomatoes as specified below under "Application Instructions".

For effective control of grasses and broadleaf weeds with postemergence applications, apply Sencor® 4 before weeds are 1-inch tall. Thorough spray coverage on weed foliage is essential for adequate control with postemergence applications.


Do not use air blast or other high pressure spray equipment to make postemergence applications of Sencor® 4. Refer to the appropriate section of this label for additional information regarding spray equipment, dilution rates, mixing, sprayer cleanup, restrictions, container disposal and cautions.

For specific application information see the "General Information" section in the front of this label.



| WEEDS CONTROLLED | | |
|---|--|---|
| PREPLANT INCORPORATED APPLICATIONS TRANSPLANT TOMATOES ONLY | | |
| Broadcast Sprays - 1/2 to 1 Pt Sencor® 4/Acre | | |
| Broadleaves Galinsoga (<i>Galinsoga</i> spp.) Lambsquarters (<i>Chenopodium album</i>) | *Pigweed, Redroot (<i>Amaranthus retroflexus</i>) | *Purslane, Common (<i>Portulaca oleracea</i>) |
| Grasses *Goosegrass (<i>Eleusine indica</i>) | | |
| Preplant incorporated applications applied as directed will suppress foxtails, panicums and barnyardgrass. | | |
| Sencor®/Trifluralin Tank Mix: This tank mix combination applied preplant incorporated as directed on this label will control the weeds listed above plus those weeds listed on the Trifluralin label. | | |
| * For optimum control of these weeds, use the highest rate specified on the label for the type of application to be made. Repeat postemergence applications may be needed for best control. | | |

| WEEDS CONTROLLED | | |
|---|--|--|
| POSTEMERGENCE APPLICATIONS ESTABLISHED TOMATOES | | |
| For effective control of weeds with postemergence applications, apply Sencor® 4 before weeds are 1-inch tall. | | |
| Broadcast Sprays - 1/2 to 1 Pt Sencor® 4/Acre | | |
| Broadleaves Carpetweed (<i>Mollugo verticillata</i>) Fumitory (<i>Fumaria officinalis</i>) Galinsoga (<i>Galinsoga</i> spp.) *Jimsonweed (<i>Datura stramonium</i>) | *Ladysthumb (<i>Polygonum persicaria</i>) Lambsquarters (<i>Chenopodium album</i>) Mustard, Wild (<i>Brassica kaber</i>) Pigweeds (<i>Amaranthus</i> spp.) Purslane (<i>Portulaca oleracea</i>) | *Ragweed, Common (<i>Ambrosia artemisiifolia</i>) *Smartweed, Pennsylvania (<i>Polygonum pensylvanicum</i>) Toadflax (<i>Linaria</i> spp.) *Velvetleaf (<i>Abutilon theophrasti</i>) |
| Directed Sprays -1 to 2 Pt Sencor® 4/Acre | | |
| Grasses *Foxtail, Yellow (<i>Setaria glauca</i>) | Goosegrass (<i>Eleusine indica</i>) | Plus Weeds Listed Under Broadcast Sprays |
| * For optimum control of these weeds, use the highest rate specified on the label for the type of application to be made. Repeat postemergence applications may be needed for best control. | | |
| Postemergence applications as directed on this label will suppress barnyardgrass and crabgrass when these weeds are less than 1-inch tall. | | |



| BROADCAST APPLICATION INSTRUCTIONS FOR TOMATOES | |
|---|---|
| Sencor® 4 *Pt/Acre | REMARKS |
| 1/2 to 1 | PREPLANT INCORPORATED--TRANSPLANT TOMATOES ONLY: Apply specified dosage in 10 or more gallons of water per acre as a broadcast spray to the soil surface immediately before transplanting. Incorporate to a depth of 2 to 4 inches with equipment capable of uniformly mixing the chemical into the soil. This application may be made alone or in a tank mix combination with Trifluralin E.C. When transplanting tomatoes, place the root system of the plants below the herbicide incorporation zone or injury may occur. Refer to the Trifluralin label for specific rate of application and for additional precautions and restrictions for tomatoes. |
| 1/2 to 1 | POSTEMERGENCE BROADCAST SPRAY--ESTABLISHED TOMATOES: Apply specified dosage in 20 or more gallons of water per acre as a broadcast spray, or apply in 1/4 to 3/4 inch of water (use 1/4 to 1/2 inch of water on sandy soils) per acre as a continuous injection in center pivot and lateral move systems or apply in the last 15 to 30 minutes of set in permanent solid set sprinkler systems. One or more applications may be applied per use season. Allow at least 14 days between applications or severe crop injury may occur. For transplanted tomatoes, do not apply until transplants have recovered from transplant shock and new growth is evident. Do not apply to tomatoes within 24 hours of application of other pesticides. Do not tank mix with other pesticides. (See "Special Precautions" below.) |
| 1 to 2 | POSTEMERGENCE DIRECTED SPRAY--ESTABLISHED TOMATOES: Apply specified dosage in 20 or more gallons of water per acre as a directed spray. One or more applications may be applied per use season. Allow at least 14 days between applications or severe crop injury may occur. Avoid contacting tomato foliage with spray. This method of treatment is recommended for use in fields with a history of severe weed pressure or in fields infested with hard-to-control weeds. For transplanted tomatoes, do not apply until transplants have recovered from transplant shock and new growth is evident. Do not apply to tomatoes within 24 hours of application of other pesticides. (See "Special Precautions" below.) When banding see the appropriate section in the front of this label. |
| * Use the higher rate in fields with a history of severe weed pressure and for maximum residual weed control. | |

SPECIAL PRECAUTIONS (Tomatoes): Do not apply more than a total of 2 pts. Sencor® 4 per crop season. Do not apply the total amount of 2 pt Sencor® 4 within a time span of less than 35 days, except in the case of directed sprays. Allow at least 14 days between applications, regardless of dosage or method of application or severe crop injury may occur. Do not apply within 7 days of harvest. Do not apply within 3 days after periods of cool, wet or cloudy weather, or crop injury will occur. Do not use hot caps on tomatoes within 7 days before or at any time after application of Sencor® 4. Do not treat seeded tomatoes until plants have reached the 5 to 6 leaf stage or severe crop injury may occur. Crop injury or delayed maturity may result from broadcast or directed spray applications if tomatoes are growing under stress conditions such as periods of drought or cool, wet and cloudy weather preceding application. For newly introduced tomato varieties with unknown tolerance to Sencor®, treat only a small area to determine if Sencor® 4 can be used without injury to the crop.

Aerial application is prohibited.

DO NOT USE SENCOR® 4 ON TOMATOES IN KERN COUNTY, CALIFORNIA.

Spring and Winter Barley and Winter Wheat

Sencor® 4 herbicide is recommended for control or suppression of certain grasses and broadleaf weeds when applied postemergence to spring and winter barley or winter wheat. Sencor® 4 alone and several tank mixture treatments are recommended for use in the following states: AR, GA, ID, IL, IN, KS, KY, LA, MS, MO, MT, NV, OH, OK, OR, TN, TX, UT, WA.

Mixing: See the "General Information" section of this label for specific mixing procedures. When tank mixing, carefully follow the instructions on this label. Refer to the other product labels registered for use on barley and winter wheat for additional use directions, rates, weeds controlled and restrictions.

Application: Sencor® 4 may be applied by aerial or ground application equipment. Use a minimum spray volume of 2 gpa by air and 10 gpa by ground. Uniform spray coverage is necessary to obtain optimum weed control and to minimize potential for crop injury. Do not exceed rates specified on this label. Do not apply Sencor® 4 through any type of irrigation equipment. Apply Sencor® 4 when the crop is healthy and actively growing. Sencor® 4 may be applied more than once per crop season. Allow a minimum of 21 days between applications if wheat is actively growing or allow 45 days between applications if wheat is growing in adverse conditions, has entered dormancy or is stressed due to frost damage, disease, drought or excessive moisture. Do not use on soils containing less than 0.75% organic matter. Do not apply more than a total of 16 fluid ounces Sencor® 4 (8 ounces active ingredient) per acre per year. On

irrigated wheat and barley, do not apply more than 0.5 inch of water for the first irrigation, the maximum amount for each additional irrigation should not exceed 1 inch. Allow a minimum of 14 days between the first irrigation and subsequent irrigations.

Performance Factors: Weed control may not be observed for 2 to 4 weeks under normal growth conditions and for 4 to 6 weeks under very dry conditions. Moisture (at least 1/2 inch) is required within 2 to 3 weeks after application to move Sencor® into the weed root zone. Lack of adequate moisture after application may result in poor or erratic weed control. Control or suppression of listed weeds is dependent on weed size at time of application. Control or suppression may be reduced if broadleaf weeds are taller than 1 inch or grasses have more than 2 leaves.


Tank Mixtures: Sencor® 4 may be tank mixed with Ally, Amber, Finesse, Glean FC, Harmony Extra, 2,4-D, MCPA, Igran, Banvel/Banvel SGF, Bronate or Buctril herbicides. A nonionic surfactant containing at least 80% active ingredient may be used in Sencor® 4 tank mixes with sulfonylurea herbicides (Ally, Amber, Finesse, Glean FC and Harmony Extra). Do **not** use a crop oil concentrate or any adjuvant containing vegetable or petroleum oils with any Sencor® 4 mix as crop injury may result. Additional pesticides may also be tank mixed with Sencor® 4 unless specifically prohibited on the mix products' label. In some instances, combinations with organophosphate insecticides may cause temporary leaf yellowing and/or crop injury, especially when widely fluctuating day/night temperatures occur near application. Always refer to the other product labels registered for use on wheat and barley for additional directions, rates and weed species controlled. Observe all precautions and limitations on labeling of all products used in mixtures.

Feeding Restrictions: Do not graze wheat within 14 days of Sencor® 4 application or harvest grain within 21 days after last application. Do not graze or harvest barley before crop maturity. For tank mix combinations, follow the most restrictive label.

Special Precautions: Wheat and Barley Injury - Crop injury may occur if Sencor® 4 is applied:

- When the crop is under stress such as winter kill, frost damage, disease, drought or excessive moisture, severe grazing, or when these conditions follow the application.
- In combination with fluid fertilizer especially with the addition of surfactant,
- Prior to the growth stage specified on this label.
- To soils high in lime or sodium, a pH greater than 7.7, calcareous, gravelly, thinly covered or exposed subsoil areas.
- To fields where wheat and barley seeds have been planted less than 1 inch deep.
- To a non-winter hardy wheat or barley variety.
- To a sensitive wheat or barley variety as listed below.
- To frozen soil or crop still in winter dormancy.

Wheat and Barley Rotations Following Potatoes Treated with Sencor® 4: If planting a sensitive wheat and barley variety (listed under the wheat and barley variety tolerance portion of this label), following potatoes treated with Sencor® 4 or metribuzin containing products, refer to the potato section of the Sencor® 4 label for special cultural practices to follow.



APPLICATION INSTRUCTIONS

Sencor® 4 alone or in a tank mix with labeled broadleaf herbicides may be applied by aerial or ground spray equipment as a broadcast postemergence spray.

| POSTEMERGENCE BROADCAST APPLICATION INSTRUCTIONS FOR SENCOR® 4 | | | |
|--|--------------|--------------------------|------------|
| CROP GROWTH STAGE | SOIL TEXTURE | SENCOR® 4 RATE (fl oz/A) | |
| | | % ORGANIC MATTER | |
| | | 0.75 to 2.0 | OVER 2.0 |
| 2 Leaf To 2 Tiller | Coarse | 1.5 to 3 | 1.5 to 4.5 |
| | Medium | 1.5 to 4.5 | 3 to 4.5 |
| | Fine | 3 to 4.5 | 3 to 6 |
| Use these rates on crops with secondary roots smaller than 1 inch. For dryland winter wheat (nonirrigated), apply the highest specified rate to achieve maximum weed suppression/control | | | |
| 3 Tiller To 4 Tiller | Coarse | 4.5 to 6 | 6 to 7.5 |
| | Medium | 6 to 7.5 | 7.5 to 9 |
| | Fine | 7.5 to 9 | 7.5 to 9 |
| Do not apply within 2 weeks after grazing or breaking of winter dormancy. Apply after the crop is at or beyond the 3 tiller growth stage but before jointing. Secondary roots should be developed and larger than 1 inch long. Do not apply before 75 days after planting. For dryland winter wheat (nonirrigated), apply the highest specified rate to achieve maximum weed suppression/control. GEORGIA ONLY: Wheat must be planted before November 15 in the Piedmont area and Northern part of the state, and before December 1 in the Coastal Plain area. | | | |
| Over 4 Tillers | Coarse | 6 to 9 | 7.5 to 12 |
| | Medium | 6 to 12 | 7.5 to 12 |
| | Fine | 7.5 to 12 | 12 to 16 |
| Do not apply within 2 weeks after grazing or breaking of winter dormancy. Apply after the crop is at or beyond the 3 tiller growth stage but before jointing. Secondary roots should be developed and larger than 1 inch long. Do not apply before 75 days after planting. For dryland winter wheat (nonirrigated), apply the highest specified rate to achieve maximum weed suppression/control. GEORGIA ONLY: Wheat must be planted before November 15 in the Piedmont area and Northern part of the state, and before December 1 in the Coastal Plain area. | | | |

WHEAT AND BARLEY VARIETAL TOLERANCE*

Wheat and barley varieties vary in their tolerance to Sencor® 4. Varieties below are tolerant to and are recommended for use with Sencor® 4:

Winter Wheat: Abe, AgriPro Mason, AgriPro Shiloh, Arthur, AS 7846, AS 7853, Baker Seed 32, Barbie VI, Basin, Batum, Bayies, Becker, Bintee V, Buchshot DS 2368, Caldwell, Cardinal, Cashup, Centurk, Cherokee, Cheyenne, Clark, Coker 747, Coker 762, Coker 797, Coker 88-15, Coker 9134, Coker 9543, Coker 9904, Coker 9907, Daws, DB 533W, DB 562W, DB 580W, Delta King 502, Delta King 9027, Dixie 952, Doublecrop, Dusty, Dyna-gro 426, Dynasty, Excel, Faro, FFR 525W, Florida 302, FS 432, FS 433, FS 435, Gains, Garst 64, Georgia 100, Genie V, Hatton, Hawk, Hill 81, Howell, Hunter, Hyak, Hyslop, Katie VI, KY 16-2, Larned, Lewis 833, Lewjain, Lisa, Longhorn, Luke, Madison, Magnum, Malcom, McDermid, McNair 1003, McNair 1813, Molly, Moro, Naely, Nelson, Newton, Norstar, Norwin, Nugaines, Oasis, Omega 78, Paha, Peck, Pike, Pioneer 2157, Pioneer 2180, Pioneer 2510, Pioneer 2545, Pioneer 2548, Pioneer 2550, Pioneer 2552, Pioneer 2555, Pioneer 2566, Pioneer 2571, Pioneer 2580, Pioneer 2684, Quantum 577, Redwin, Rocky, Saluda, Sawyer, SC 104, Siouxland, Sprague, Southern Belle, Stacy, Stallion, Stephens, TAM W101, TAM 105, TE 377, TE 2548, TE SR204, TR 8555, TR 8557, TR 8768, Tiber, Tomahawk, Traveler, Tres, Tyee, Tyler, Verne, Victory Wakefield, Wahser, Weston, Winalta, Wrangler.

Barley: Advance, Boyer, Clark, Compana, Hannchen, Hector, Hesk, Hudson, Lud, Luther, Kamiak, Klages, Olympic, Piro, Steptoe and Triumph.

Do not use the following wheat and barley varieties which are sensitive to Sencor® 4:

Winter Wheat: AgriPro Clemens, AT 90W, AT 91W, Arapaho, Baker Seed 33, Century, Cimarron, Coker 833, Coker 916, Coker 983, Coker 9024, Coker 9105, Coker 9323, Coker 9663, Choker 9474, Choker 9835, Choker 9766, Choker 9877, EK 102, EK 114, FAR 555, Florida 304, Freedom, FS 417, FS 423, FS 425, FS 430, Gore, Hazen, Hickory, Jackson, Julie III, KY-49-25, Linden, Madison, Mesa,

Mustang, Pacer, Pioneer 2551, Pioneer 2163, Pioneer 2643, Pioneer 2691, Princeton 733, PER W71, PER 226, PER 278, Rosen, Savannah, Sierra, TAM 107, TR 101, TR 1011, TR 8822, Triumph 64, Vona, Wings, Winridge, Yamhill.

Spring/Durum Wheat: Avoid use on Spring wheat and Durum wheat varieties.

Barley: Glenn, Morex, Moravian 3, Larker, Summit, Bracken, Anheuser Busch B2601 and varieties with Morex parentage.

Varieties Not Listed: To avoid possible crop injury on any variety not mentioned in this label, contact a Bayer CropScience representative or herbicide expert for a variety recommendation prior to treatment or treat a small strip of the unlisted variety with the specified Sencor® 4 rate to ascertain crop tolerance before treating an entire field.

* Abbreviated names of vendors: AS (Agseco), AT (Agratech), DB (Diener Bros.), FS (Growmark FS), PI (Pioneer), PER (Hybritech), SC (J.M. Schultz), TE (Terra), and TR (Terral).

Weeds Controlled

Used at specified rates, Sencor® 4 will control many annual broadleaf weeds. Control is best when applied to young, actively growing weeds. Weeds controlled by Sencor® 4 include:

| | | |
|---------------------------|----------------------|-------------------------------------|
| Bittercress | Filaree, Redstem | Pepperweed, Virginia |
| Catchfly, Conical (Sand) | Geranium, Carolina | Pigweed, spp. |
| Catchweed (Madwort) | Gromwell, spp. | Pineappleweed |
| Chickweed, Common | Henbit | Polemonium, Annual (Jacob's Ladder) |
| Chickweed, Mouseear | Knotweed, Prostrate | Radish, Wild |
| Corncockle | Lambsquarter, Common | Shepardspurse |
| Dogfennel (Mayweed) | Lettuce, Miners | Speedwell, Ivyleaf |
| Evening Primrose, Cutleaf | Mustard, Blue | Turnip, Wild |
| Falseflax, Smallseed | Mustard, Wild | |
| Fiddleneck, Tarweed | Pennycress, Field | |

Weeds Suppressed

Sencor® 4 control of the following weeds varies from poor to excellent depending on time of application, stage of growth at application, temperatures and soil moisture conditions following treatment. For maximum effect on these weeds, apply the highest specified rate at the earliest growth stage timing for each particular soil type and organic matter. Suppression is a reduction in weed size and growth as compared to a non-treated area in the same field.

| | | |
|--|-----------------------------|-------------------------------|
| BROADLEAVES | | |
| Buckwheat, Wild* | Kochia* | Tanysmustard |
| Buttercup, spp. | Lettuce, Prickly | Thistle, Russian |
| Cowcockle | Mustard, Tumble (Jim Hill)* | Vetch, Winter |
| GRASSES | | |
| Barley, Hare (Wild) | Brome, Downy* | Oat, Wild* |
| Barley, Little | Brome, Japanese* | Rescuegrass* |
| Blackgrass | Brome, Ripgut* | Whitlowgrass, Spring (Vernal) |
| Bluegrass, Annual | Cheat* | Windgrass |
| Bluegrass, Bulbous | Foxtail, spp* | |
| * Use the highest specified Sencor® 4 rate for maximum weed suppression. | | |

FOR WEED CONTROL IN A WHEAT/FALLOW/WHEAT ROTATION (Idaho, Oregon, Utah and Washington Only)

Sencor® 4 may be applied to provide weed control during the fallow period after wheat harvest or in the Spring before winter wheat is planted. Winter wheat can be seeded 4 months (120 days) after Spring application. Mechanical tillage or the application of a contact herbicide may be required to control weeds germinating prior to seeding of winter wheat. Best results will be obtained where straw and chaff are evenly distributed across the field.

For specific application information see the "General Information" section in the front of this label.

Where weed growth is present at application time, Sencor® 4 should be applied with Gramoxone or other contact herbicide. Refer to the other product label registered for additional directions, rates, and weed species controlled.

Weeds Controlled

| BROADLEAVES | | |
|--|--|--|
| Chickweed, Common (<i>Stellaria media</i>) | Mustard, Jim Hill (<i>Sisymbrium altissimum</i>) | Pigweeds (<i>Amaranthus</i> spp.) *Russian thistle (<i>Salsola iberica</i>) *Wild Sunflower (<i>Helianthus</i> spp.) |
| Henbit (<i>Lamium amplexicaule</i>) | Mustard, Tansy (<i>Descurainia pinnata</i>) | |
| *Kochia (<i>Kochia scoparia</i>) | Mustard, Treacle (<i>Eysimum repandum</i>) | |
| Lambsquarters (<i>Chenopodium album</i>) | Mustard, Wild (<i>Brassica kaber</i>) | |
| Mustard, Blue or Purple (<i>Chorispora tenella</i>) | Pennycress, Field (Fanweed) (<i>Thlaspi arvense</i>) | |
| GRASSES | | |
| Cheatgrass (<i>Bromus secalinus</i>) | Downy Brome (<i>Bromus tectorum</i>) | *Wheat, Volunteer (<i>Triticum</i> spp.) |
| * Note: Since control of these weeds may be variable depending on moisture following application, the higher labelled rate is recommended. | | |

After Harvest Application (Fall Fallow): Sencor® 4 may be applied to wheat stubble after harvest in the Fall. Apply 1 to 1-1/4 pt per acre broadcast before weeds emerge. Use higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

Do not plant crops in treated areas for at least 10 months following Fall applications. Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.

Sencor® 4 may be applied at 1 to 1-1/4 pt per acre as directed above for a Fall application. If other vegetation is present at the time of application use a contact herbicide.

Spring Application (Summer Fallow): Sencor® 4 may be applied to wheat stubble in the Spring. Apply 3/4 to 1 pt per acre broadcast before weeds emerge in the Spring. Use higher rate for longer weed control or weeds designated as requiring higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

Precautions and Restrictions: Do not graze treated fields. Do not plant Spring seeded barley following Fall fallow applications of Sencor®. Where Sencor® was applied in the Fall, do not apply Sencor® in the Spring.

FOR WEED CONTROL IN A FALLOW ROTATION WITH BARLEY AND WHEAT

(Colorado, Kansas, Montana, Nebraska, and Wyoming Only)

Sencor® 4 may be applied to provide weed control during the fallow period after wheat or barley harvest or in the Spring before planting of Winter wheat or barley. Mechanical tillage or the application of a contact herbicide may be required to control weeds germinating prior to seeding of Winter wheat or barley.

For specific application information see the "General Information" section in the front of this label.

Where weed growth is present at application time, Sencor® 4 should be applied with Gramoxone, Roundup, or other contact herbicide. Refer to the other product label registered for additional directions, rates, and weed species controlled. Do not plant crops in treated areas earlier than 10 months following Fall applications. Do not rotate any crop not listed on this label for 18 months following application of Sencor® 4.

Weeds Controlled

| BROADLEAVES | | |
|--|---|--|
| Chickweed, Common (<i>Stellaria media</i>) | Mustard, Blue or Purple (<i>Chorispora tenella</i>) | Pennycress, Field (fanweed) (<i>Thlaspi arvense</i>) |
| Cowcockle (<i>Vaccaria pyramidata</i>) | Mustard, Jim Hill (<i>Sisymbrium altissimum</i>) | Pigweeds (<i>Amaranthua</i> spp.) |
| Henbit (<i>Lamium amplexicaule</i>) | Mustard, Tansy (<i>Descurainia pinnata</i>) | Russian thistle (<i>Salsola iberica</i>) |
| *Kochia (<i>Kochia scoparia</i>) | Mustard, Treacle (<i>Eysimum repandum</i>) | Sunflower (<i>Helianthus</i> spp.) |
| Lambsquarters (<i>Chenopodium album</i>) | Mustard, Wild (<i>Brassica kaber</i>) | |
| GRASSES | | |
| Cheatgrass (<i>Bromus secalinus</i>) | *Foxtail, Green (<i>Setaria viridis</i>) | *Wild Oats (<i>Avena fatua</i>) |
| Downy Brome (<i>Bromus tectorum</i>) | *Wheat, Volunteer (<i>Triticum</i> spp.) | |
| *Note: Since control of these weeds may vary depending on moisture following application, use the higher rate specified below. | | |

AFTER HARVEST APPLICATION (Fall Fallow): Sencor® 4 may be applied to the stubble after harvest in the Fall. Apply 1-1/4 to 1-1/2 pt per acre broadcast before weeds emerge. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

SPRING APPLICATION (Summer Fallow): Sencor® 4 may be applied to the stubble in the Spring. Apply 3/4 to 1 pt per acre broadcast before weeds emerge in the Spring. Use the higher rate for longer weed control or weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation. Wheat or barley can be seeded 120 days after Spring application.

Precautions and Restrictions: Do not graze treated fields. Do not plant Spring seeded barley following Fall applications for fallow. Where Sencor® was applied in the Fall, do not apply Sencor® in the Spring.

CROP ROTATION RESTRICTIONS

Waiting Period After Sencor® 4 Flowable Herbicide Application¹


| | | | |
|------------------|--|--|---|
| 4 Months | Alfalfa Asparagus Barley ² Corn | Forage Grasses Sainfoil Soybeans | Sugarcane Tomatoes Wheat ² |
| 8 Months | Barley Lentils | Peas Wheat | |
| 12 Months | Potatoes | Rice ³ | |
| 18 Months | Sugar Beets, Onions and other root crops not listed on this label, and all other crops not listed on this label. | | |

¹ Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed. Stand reductions may occur in some areas.

² Following peas, lentils or soybeans.

³ Do not rotate rice after any application to a primary crop greater than 1.0 lb ai/A of Sencor® 4 per season.

Do not rotate any crop not listed on this label after application of Sencor® 4 to sugarcane.



IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.


NET CONTENTS:

Sencor, Bronate and Buctril are registered trademarks of Bayer.
Banvel is a registered trademark of Velsicol Chemical Corp.
Command and Commence are registered trademarks of FMC Corp.
Amber and Igran are registered trademarks of Ciba-Geigy.
Ally, Bladex, Canopy, Finesse, Glean and Harmony are registered trademarks of E. I. duPont de Nemours and Co.
Gramoxone is a registered trademark of ICI Americas, Inc.
Freedom and Lasso are registered trademarks of Monsanto Co.
Roundup is a trademark of Monsanto Co.
Pursuit, Prowl and Scepter are registered trademarks of American Cyanamid Co.
Sonalan and Treflan are trademarks of DowElanco Products Co.
X-77 is a registered trademark of Valent USA Corp.

Produced for



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)



Sencor 4 Flowable Herbicide (PENDING) 05/24/10, Resubmitted 06/01/10